

## Systematic Review

# A Systematic Review and Best Evidence Synthesis of Effectiveness of Therapeutic Facet Joint Interventions in Managing Chronic Spinal Pain

Laxmaiah Manchikanti, MD<sup>1</sup>, Alan D. Kaye, MD, PhD<sup>2</sup>, Mark V. Boswell, MD, PhD<sup>3</sup>, Sanjay Bakshi, MD<sup>4</sup>, Christopher G. Gharibo, MD<sup>5</sup>, Vahid Grami, MD<sup>6</sup>, Jay S. Grider, DO, PhD<sup>7</sup>, Sanjeeva Gupta, MD<sup>8</sup>, Sachin "Sunny" Jha, MD<sup>9</sup>, Dharam P. Mann, MD<sup>10</sup>, Devi E. Nampiaparampil, MD<sup>11</sup>, Manohar Lal Sharma, MD<sup>12</sup>, Lindsay N. Shroyer, MD<sup>13</sup>, Vijay Singh, MD<sup>14</sup>, Amol Soin, MD<sup>15</sup>, Ricardo Vallejo, MD<sup>16</sup>, Bradley W. Wargo, DO<sup>17</sup>, and Joshua A. Hirsch, MD<sup>18</sup>

From: <sup>1</sup>Pain Management Center of Paducah, Paducah, KY, and University of Louisville, Louisville, KY; <sup>2</sup>LSU Health Science Center, New Orleans, LA; <sup>3</sup>Department of Anesthesiology and Perioperative Medicine, University of Louisville, Louisville, KY; <sup>4</sup>Manhattan Spine and Pain Medicine, New York, NY; <sup>5</sup>NYU Langone - Hospital for Joint Diseases, NYU School of Medicine, New York, NY; <sup>6</sup>Geisinger Medical Center Interventional Pain Center Woodbine, Danville, PA; <sup>7</sup>UK Healthcare Pain Services, University of Kentucky, Lexington, KY; <sup>8</sup>Bradford Teaching Hospitals NHS Foundation Trust, Bradford, UK; <sup>9</sup>Rush University Medical Center (RUMC), Chicago, IL; <sup>10</sup>Garden State Pain Management, Whiting, NJ; <sup>11</sup>New York University School of Medicine, New York, NY; <sup>12</sup>The Walton Centre for Neurology and Neurosurgery NHS Foundation Trust, Liverpool, UK; <sup>13</sup>Florida Pain Management Associates, Sebastian, FL; <sup>14</sup>Spine Pain Diagnostics Associates, Niagara, WI; <sup>15</sup>Ohio Pain Clinic, Centerville, OH; <sup>16</sup>Millennium Pain Center, Bloomington, IL; <sup>17</sup>Illinois State University, Normal, IL; <sup>18</sup>The McFarland Clinic, Mary Greeley Medical Center, Department of Pain Medicine, Ames, IA; <sup>19</sup>Massachusetts General Hospital and Harvard Medical School, Boston, MA.

Address Correspondence:  
Laxmaiah Manchikanti, M.D.  
2831 Lone Oak Road  
Paducah, Kentucky 42003  
E-mail: drlm@thepainmd.com :

Author Affiliation and Disclaimers  
on P. E573-E574

Manuscript received: 06-14-2015  
Accepted for publication:  
07-09-2015

Free full manuscript:  
www.painphysicianjournal.com

**Background:** The therapeutic spinal facet joint interventions generally used for the treatment of axial spinal pain of facet joint origin are intraarticular facet joint injections, facet joint nerve blocks, and radiofrequency neurotomy. Despite interventional procedures being common as treatment strategies for facet joint pathology, there is a paucity of literature investigating these therapeutic approaches.

Systematic reviews assessing the effectiveness of various therapeutic facet joint interventions have shown there to be variable evidence based on the region and the modality of treatment utilized. Overall, the evidence ranges from limited to moderate.

**Objective:** To evaluate and update the clinical utility of therapeutic lumbar, cervical, and thoracic facet joint interventions in managing chronic spinal pain.

**Study Design:** A systematic review of therapeutic lumbar, cervical, and thoracic facet joint interventions for the treatment of chronic spinal pain.

**Methods:** The available literature on lumbar, cervical, and thoracic facet joint interventions in managing chronic spinal pain was reviewed. The quality assessment criteria utilized were the Cochrane Musculoskeletal Review Group criteria and Interventional Pain Management Techniques – Quality Appraisal of Reliability and Risk of Bias Assessment (IPM – QRB) for randomized trials and Interventional Pain Management Techniques – Quality Appraisal of Reliability and Risk of Bias Assessment for Nonrandomized Studies (IPM – QRBNR) for observational studies.

The level of evidence was classified at 5 levels from Level I to Level V.

Data sources included relevant literature identified through searches on PubMed and EMBASE from 1966 through March 2015, and manual searches of the bibliographies of known primary and review articles.

**Outcome Measures:** The primary outcome measure was pain relief (short-term relief = up to 6 months and long-term > 6 months). Secondary outcome measures were improvement in functional status, psychological status, return to work, and reduction in opioid intake consumption.

**Results:** A total of 21 randomized controlled trials meeting appropriate inclusion criteria were assessed in this evaluation. A total of 5 observational studies were assessed.

In the lumbar spine, for long-term effectiveness, there is Level II evidence for radiofrequency neurotomy and lumbar facet joint nerve blocks, whereas the evidence is Level III for lumbosacral intraarticular injections.

In the cervical spine, for long-term improvement, there is Level II evidence for cervical radiofrequency neurotomy and cervical facet joint nerve blocks, and Level IV evidence for cervical intraarticular injections.

In the thoracic spine there is Level II evidence for thoracic facet joint nerve blocks and Level IV evidence for radiofrequency neurotomy for long-term improvement.

**Limitations:** The limitations of this systematic review include an overall paucity of high quality studies and more specifically the lack of investigations related to thoracic facet joint injections.

**Conclusion:** Based on the present assessment for the management of spinal facet joint pain, the evidence for long-term improvement is Level II for lumbar and cervical radiofrequency neurotomy, and therapeutic facet joint nerve blocks in the cervical, thoracic, and lumbar spine; Level III for lumbar intraarticular injections; and Level IV for cervical intraarticular injections and thoracic radiofrequency neurotomy.

**Key Words:** Spinal pain, chronic low back pain, chronic neck pain, chronic thoracic pain, intraarticular facet joint blocks, facet joint nerve blocks, conventional radiofrequency neurotomy, pulsed radiofrequency neurolysis

### **Pain Physician 2015; 18:E535-E582**

---

**C**hronic spinal pain with or without extremity pain, chest wall pain, or headaches is commonly encountered in modern day health care, at a time when health care costs and disability are exploding with a corresponding exponential increase in treatment modalities (1-15). Controlled studies have previously established intervertebral discs, facet joints, and sacroiliac joints as potential sources of spinal and extremity pain (10,16-20). It has been described that facet joint degeneration can result from abnormal motion associated with disc degeneration, as well as arthritis, similar to that seen in other synovial joints (21-24). In addition, multiple mechanisms have been postulated as being responsible for modulation of spinal pain related to the facet joints, including capsular stretch, entrapment of synovial villi between the articular surfaces, nerve impingement by osteophytes, and release of inflammatory substances (25-29). The spinal facet joints have been shown to have an abundant nerve supply (30-41); they can cause pain much like what has been reported in normal volunteers who have persistent spinal pain and referred pain into the extremities, chest wall, or head (42-53); are known to be susceptible to arthritic changes, degenerative changes, inflammation, and injury, all of which can lead to a restriction in range of motion and pain upon movement (21-29). By using accepted and proven diagnostic techniques, facet joints have been shown to be a pain generator with subsequent therapeutic application of modalities with significant improvement in pain and disability directed at facet joint innervation (10,16-19,54). Thus, facet joint pain may be diagnosed with reliability based on established controlled diagnostic blocks and may be managed with therapeutic interventions including intraarticular injections, facet joint nerve blocks, or facet joint neurolytic procedures (10,55-57). The diagnostic accuracy and reliability of facet joint nerve blocks for chronic spinal pain have

shown that with controlled diagnostic blocks, there is a prevalence of 27% to 41% in the low back with a false-positive rate of 25% to 44%, a prevalence of 36% to 67% and a false-positive rate of 27% to 63% in the cervical spine, and finally, in the thoracic spine a prevalence rate of 34% to 48% with a false-positive rate of 42% to 48% (19).

The accurate selection of patients for therapeutic modalities and diagnostic accuracy is crucial, since interventional techniques, specifically facet joint interventions, have shown overall increases of 293% or 11.1% per year per 100,000 fee-for-service Medicare population from 2000 to 2013, compared to 14% of US population and 64% of Medicare beneficiaries (13-15). In fact, lumbar facet joint nerve blocks have increased at a rate of 213%, cervical/thoracic facet joint nerve blocks have increased at a rate of 350%, and radiofrequency neurotomy have increased even more at a rate of 522% for lumbar facet joint radiofrequency neurotomy and 845% for cervical facet joint radiofrequency neurotomy from 2000 to 2013 in the fee-for-service Medicare population (13-15). Based on the selection criteria of appropriate diagnosis with controlled diagnostic blocks, multiple therapeutic interventions have been assessed in multiple reviews (10,55-61). These systematic reviews demonstrated variable evidence for facet joint neurotomy, facet joint nerve blocks, and intraarticular injections (55-61).

Thus, the debate over the appropriateness of these procedures, continues often with contradictory evidence based on personal and/or professional bias, conservatism, and policy implementations, but not grounded in an appropriate synthesis of the literature (7,58-68). Multiple systematic reviews have been performed based on methodologic assessment but we believe that they display significant bias and have made methodologic errors.

The objective of this systematic review is to assess

the effectiveness of therapeutic facet joint interventions with appropriate methodology.

## 1.0 METHODS

The methodology utilized in this systematic review followed the review process derived from evidence-based systematic reviews and meta-analyses of randomized trials and observational studies (69-73).

### 1.1 Criteria for Considering Studies for This Review

#### 1.1.1 Types of Studies

- Randomized controlled trials
- Observational studies

#### 1.1.2 Types of Participants

Patients suffering with chronic neck pain, mid back pain, upper back pain or low back pain of at least 3 months duration.

Patients with acute trauma, fractures, malignancies, and inflammatory diseases were excluded.

#### 1.1.3 Types of Interventions

Cervical, thoracic, and lumbar facet joint interventions appropriately performed with proper technique under image guidance (fluoroscopy, computed tomography [CT], or magnetic resonance imaging [MRI]) were included. Blind and ultrasound-guided interventions were excluded.

#### 1.1.4 Types of Outcome Measures

- ◆ The primary outcome parameter was pain relief.
- ◆ The secondary outcome measure was functional status improvement.

### 1.2 Literature Search

All of the available trials in all languages from all countries providing appropriate management with outcome evaluations were considered for inclusion. Searches were performed from the following sources without language restrictions:

1. PubMed from 1966  
[www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed](http://www.ncbi.nlm.nih.gov/sites/entrez?db=pubmed)
2. Cochrane Library  
[www.thecochranelibrary.com/view/0/index.html](http://www.thecochranelibrary.com/view/0/index.html)
3. U.S. National Guideline Clearinghouse (NGC)  
[www.guideline.gov/](http://www.guideline.gov/)
4. Previous systematic reviews and cross references

5. Clinical Trials  
[clinicaltrials.gov/](http://clinicaltrials.gov/)
6. All other sources including non-indexed journals and abstracts

The search period was from 1966 through March 2015.

### 1.3 Search Strategy

The search strategy emphasized chronic spinal pain treated with facet joint interventions. The search terms included: cervical, mid back, and low back pain, facet or zygapophysial joint pain, cervical, thoracic, and lumbar facet injections; cervical, thoracic, and lumbar facet joint nerve blocks; and cervical, thoracic, and lumbar radiofrequency thermoneurolysis.

Search criteria were as follows:

((((((((((((((((chronic low back pain) OR chronic back pain) OR chronic neck pain) OR disc herniation) OR discogenic pain) OR facet joint pain) OR herniated lumbar discs) OR nerve root compression) OR lumbosciatic pain) OR postlaminectomy) OR lumbar surgery syndrome) OR radicular pain) OR radiculitis) OR sciatica) OR spinal fibrosis) OR spinal stenosis) OR zygapophysial)) AND ((((((facet joint) OR zygapophysial) OR zygapophysial) OR medial branch block) OR diagnostic block) OR radiofrequency) OR intraarticular)

### 1.4 Data Collection and Analysis

The review focused on randomized trials for efficacy. If there were not at least 5 randomized trials, observational studies were utilized. Those patients being studied were those who had been suffering with chronic pain for 3 months or longer. Only facet joint interventions were evaluated. All of the available studies, in all languages, from all countries providing appropriate management and with outcome evaluations of 3 months or longer and statistical evaluations were reviewed. Reports without an appropriate diagnosis, nonsystematic reviews, nonrandomized studies, book chapters, and case reports were excluded.

#### 1.4.1 Inclusion and Exclusion Criteria

Randomized trials with at least 3 months of follow-up and with at least 25 patients in each group or with appropriate sample size determination were included. Observational studies of at least 50 patients with a minimum 6 month follow-up were included when needed (fewer than 5 randomized trials available in any category).

### 1.4.2 Methodological Quality or Validity Assessment

The quality of each individual article used in this analysis was assessed by Cochrane review criteria (Table 1) (70), Interventional Pain Management techniques – Quality Appraisal of Reliability and Risk of Bias Assessment (IPM – QRB) for randomized trials (Table 2) (72), and for observational studies: Interventional Pain Management Techniques – Quality Appraisal of Reliability and Risk of Bias Assessment for Nonrandomized Studies (IPM – QRBNR) (Table 3) (73).

Utilizing Cochrane review criteria, studies meeting the inclusion criteria with a score of at least 8 of 12 were considered high quality and 4 to 7 were considered moderate quality. Those with a score of less than 4 were considered low quality and were excluded.

Based on IPM-QRB criteria for randomized trials, the trials meeting the inclusion criteria that scored less than 16 were considered as low quality and were excluded, those scoring 16 to 31 were considered moderate quality, and those scoring 32 to 48 were considered high quality.

Based on IPM-QRBNR criteria for observational studies, manuscripts meeting the inclusion criteria scoring less than 16 were considered low quality and were excluded, manuscripts scoring 16 to 31 were considered moderate quality, and manuscripts scoring 32 to 48 were considered high quality.

### 1.4.3 Data Extraction and Management

Working independently and in an unblinded, standardized way, 2 review authors established the search criteria, searched for relevant literature, selected the manuscripts, and extracted the data from the included studies. Any disagreement between the 2 reviewers were discussed and debated. If no compromise was reached, another author would review the disagreement and cast the deciding opinion.

Methodological quality assessment was performed by multiple review authors with groups of 2 authors reviewing 4 to 6 manuscripts. The assessment was carried out independently in an unblinded standardized manner to assess the methodological quality and internal validity of all the studies considered for inclusion. The methodological quality assessment was performed in such a way to prevent discrepancies from occurring; if they did occur, a third reviewer was called in and the discrepancy decided by consensus. Continued issues were also discussed with the entire group and resolved.

If there was a conflict of interest with a reviewed manuscript (concerning authorship), if the reviewer was also one of the authors or there was any type of conflict, the involved authors did not review the manuscript for methodological quality assessment.

Table 1. Sources of risk of bias and Cochrane Review rating system.

A	1. Was the method of randomization adequate?	Yes/No/Unsure
B	2. Was the treatment allocation concealed?	Yes/No/Unsure
C	Was knowledge of the allocated interventions adequately prevented during the study?	
	3. Was the patient blinded to the intervention?	Yes/No/Unsure
	4. Was the care provider blinded to the intervention?	Yes/No/Unsure
	5. Was the outcome assessor blinded to the intervention?	Yes/No/Unsure
D	Were incomplete outcome data adequately addressed?	
	6. Was the drop-out rate described and acceptable?	Yes/No/Unsure
	7. Were all randomized participants analysed in the group to which they were allocated?	Yes/No/Unsure
E	8. Are reports of the study free of suggestion of selective outcome reporting?	Yes/No/Unsure
F	Other sources of potential bias:	
	9. Were the groups similar at baseline regarding the most important prognostic indicators?	Yes/No/Unsure
	10. Were co-interventions avoided or similar?	Yes/No/Unsure
	11. Was the compliance acceptable in all groups?	Yes/No/Unsure
	12. Was the timing of the outcome assessment similar in all groups?	Yes/No/Unsure

Source: Furlan AD, Pennick V, Bombardier C, van Tulder M; Editorial Board, Cochrane Back Review Group. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine (Phila Pa 1976)* 2009; 34:1929-1941 (70).

## Effectiveness of Therapeutic Facet Joint Interventions

Table 2. Item checklist for assessment of randomized controlled trials of IPM techniques utilizing IPM – QRB.

		Scoring
<b>I. TRIAL DESIGN AND GUIDANCE REPORTING</b>		
<b>1.</b>	CONSORT or SPIRIT	
	Trial designed and reported without any guidance	0
	Trial designed and reported utilizing minimum criteria other than CONSORT or SPIRIT criteria or trial was conducted prior to 2005	1
	Trial implies it was based on CONSORT or SPIRIT without clear description with moderately significant criteria for randomized trials or the trial was conducted before 2005	2
	Explicit use of CONSORT or SPIRIT with identification of criteria or trial conducted with high level reporting and criteria or conducted before 2005	3
<b>II. DESIGN FACTORS</b>		
<b>2.</b>	Type and Design of Trial	
	Poorly designed control group (quasi selection, convenient sampling)	0
	Proper active-control or sham procedure with injection of active agent	2
	Proper placebo control (no active solutions into active structures)	3
<b>3.</b>	Setting/Physician	
	General setting with no specialty affiliation and general physician	0
	Specialty of anesthesia/PMR/neurology/radiology/ortho, etc.	1
	Interventional pain management with interventional pain management physician	2
<b>4.</b>	Imaging	
	Blind procedures	0
	Ultrasound	1
	CT	2
	Fluoro	3
<b>5.</b>	Sample Size	
	Less than 50 participants in the study without appropriate sample size determination	0
	Sample size calculation with less than 25 patients in each group	1
	Appropriate sample size calculation with at least 25 patients in each group	2
	Appropriate sample size calculation with 50 patients in each group	3
<b>6.</b>	Statistical Methodology	
	None or inappropriate	0
	Appropriate	1
<b>III. PATIENT FACTORS</b>		
<b>7.</b>	Inclusiveness of Population	
<b>7a.</b>	For epidural procedures:	
	Poorly identified mixed population	0
	Clearly identified mixed population	1
	Disorders specific trials (i.e. well defined spinal stenosis and disc herniation, disorder specific, disc herniation or spinal stenosis or post surgery syndrome)	2
<b>7b.</b>	For facet or sacroiliac joint interventions:	
	No diagnostic blocks	0
	Selection with single diagnostic blocks	1
	Selection with placebo or dual diagnostic blocks	2
<b>8.</b>	Duration of Pain	
	Less than 3 months	0
	3 to 6 months	1
	> 6 months	2

Table 2 (cont.). *Item checklist for assessment of randomized controlled trials of IPM techniques utilizing IPM – QRB.*

		Scoring
<b>9.</b>	Previous Treatments	
	Conservative management including drug therapy, exercise therapy, physical therapy, etc.	
	Were not utilized	0
	Were utilized sporadically in some patients	1
	Were utilized in all patients	2
<b>10.</b>	Duration of Follow-up with Appropriate Interventions	
	Less than 3 months or 12 weeks for epidural or facet joint procedures, etc., and 6 months for intradiscal procedures and implantables	0
	3 to 6 months for epidural or facet joint procedures, etc., or 1 year for intradiscal procedures or implantables	1
	6 months to 17 months for epidurals or facet joint procedures, etc., and 2 years or longer for discal procedures and implantables	2
	18 months or longer for epidurals and facet joint procedures, etc., or 5 years or longer for discal procedures and implantables	3
<b>IV.</b>	<b>OUTCOMES</b>	
<b>11.</b>	Outcomes Assessment Criteria for Significant Improvement	
	No descriptions of outcomes OR < 20% change in pain rating or functional status	0
	Pain rating with a decrease of 2 or more points or more than 20% reduction OR functional status improvement of more than 20%	1
	Pain rating with decrease of $\geq 2$ points AND $\geq 20\%$ change or functional status improvement of $\geq 20\%$	2
	Pain rating with a decrease of 3 or more points or more than 50% reduction OR functional status improvement with a 50% or 40% reduction in disability score	2
	Significant improvement with pain and function $\geq 50\%$ or 3 points and 40% reduction in disability scores	4
12.	Analysis of All Randomized Participants in the Groups	
	Not performed	0
	Performed without intent-to-treat analysis without inclusion of all randomized participants	1
	All participants included with or without intent-to-treat analysis	2
13.	Description of Drop Out Rate	
	No description of dropouts, despite reporting of incomplete data or $\geq 20\%$ withdrawal	0
	Less than 20% withdrawal in one year in any group	1
	Less than 30% withdrawal at 2 years in any group	2
14.	Similarity of Groups at Baseline for Important Prognostic Indicators	
	Groups dissimilar with significant influence on outcomes with or without appropriate randomization and allocation	0
	Groups dissimilar without influence on outcomes despite appropriate randomization and allocation	1
	Groups similar with appropriate randomization and allocation	2
15.	Role of Co-Interventions	
	Co-interventions were provided but were not similar in the majority of participants	0
	No co-interventions or similar co-interventions were provided in the majority of the participants	1
<b>V.</b>	<b>RANDOMIZATION</b>	
16.	Method of Randomization	
	Quasi randomized or poorly randomized or not described	0
	Adequate randomization (coin toss, drawing of balls of different colors, drawing of ballots)	1

## Effectiveness of Therapeutic Facet Joint Interventions

Table 2 (cont.). *Item checklist for assessment of randomized controlled trials of IPM techniques utilizing IPM – QRB.*

	High quality randomization (computer generated random sequence, pre-ordered sealed envelopes, sequentially ordered vials, telephone call, pre-ordered list of treatment assignments, etc.)	2
<b>VI. ALLOCATION CONCEALMENT</b>		
17.	Concealed Treatment Allocation	
	Poor concealment of allocation (open enrollment) or inadequate description of concealment	0
	Concealment of allocation with borderline or good description of the process with probability of failure of concealment	1
	High quality concealment with strict controls (independent assignment without influence on the assignment sequence)	2
<b>VII. BLINDING</b>		
18.	Patient Blinding	
	Patients not blinded	0
	Patients blinded adequately	1
19.	Care Provider Blinding	
	Care provider not blinded	0
	Care provider blinded adequately	1
20.	Outcome Assessor Blinding	
	Outcome assessor not blinded or was able to identify the groups	0
	Performed by a blinded independent assessor with inability to identify the assignment-based provider intervention (i.e., subcutaneous injection, intramuscular distant injection, difference in preparation or equipment use, numbness and weakness, etc.)	1
<b>VIII. CONFLICTS OF INTEREST</b>		
21.	Funding and Sponsorship	
	Trial included industry employees	-3
	Industry employees involved; high levels of funding with remunerations by industry or an organization funded with conflicts	-3
	Industry or organizational funding with reimbursement of expenses with some involvement	0
	Industry or organization funding of expenses without involvement	1
	Funding by internal resources only with supporting entity unrelated to industry	2
	Governmental funding without conflict such as NIH, NHS, AHRQ	3
22.	Conflicts of Interest	
	None disclosed with potential implied conflict	0
	Marginally disclosed with potential conflict	1
	Well disclosed with minor conflicts	2
	Well disclosed with no conflicts	3
	Hidden conflicts with poor disclosure	-1
	Misleading disclosure with conflicts	-2
	Major impact related to conflicts	-3
<b>TOTAL MAXIMUM</b>		<b>48</b>

Source: Manchikanti L, Hirsch JA, Cohen SP, Heavner JE, Falco FJE, Diwan S, Boswell MV, Candido KD, Onyewu O, Zhu J, Sehgal N, Kaye AD, Benyamin RM, Helm II S, Singh V, Datta S, Abdi S, Christo PJ, Hameed H, Hameed M, Vallejo R, Pampati V, Racz GB, Raj PP. Assessment of methodologic quality of randomized trials of interventional techniques: Development of interventional pain management specific instrument. *Pain Physician* 2014; 17:E263-E290 (72).

Table 3. *IPM checklist for assessment of nonrandomized or observational studies of IPM techniques utilizing IPM-QRBNR.*

I. STUDY DESIGN AND GUIDANCE REPORTING		Scoring
1.	STROBE or TREND Guidance	
	Case Report/Case Series	0
	Study designed without any guidance	1
	Study designed with minimal criteria and reporting with or without guidance	2
	Study designed with moderately significant criteria or implies it was based on STROBE or TREND without clear description or the study was conducted before 2011 or similar criteria utilized with study conducted before 2011	3
	Designed with high level criteria or explicitly uses STROBE or TREND with identification of criteria or conducted prior to 2011	4
II. DESIGN FACTORS		
2.	Study Design and Type	
	Case report or series (uncontrolled – longitudinal)	0
	Retrospective cohort or cross-sectional study	1
	Prospective cohort case-control study	2
	Prospective case control study	3
	Prospective, controlled, nonrandomized	4
3.	Setting/Physician	
	General setting with no specialty affiliation and general physician	0
	Specialty of anesthesia/PMR/neurology, etc.	1
	Interventional pain management with interventional pain management physician	2
4.	Imaging	
	Blind procedures	0
	Ultrasound	1
	CT	2
	Fluoro	3
5.	Sample Size	
	Less than 100 participants without appropriate sample size determination	0
	At least 100 participants in the study without appropriate sample size determination	1
	Sample size calculation with less than 50 patients in each group	2
	Appropriate sample size calculation with at least 50 patients in each group	3
	Appropriate sample size calculation with 100 patients in each group	4
6.	Statistical Methodology	
	None	0
	Some statistics	1
	Appropriate	2
III. PATIENT FACTORS		
7.	Inclusiveness of Population	
7a.	For epidural procedures:	
	Poorly identified mixed population	1
	Poorly identified mixed population with large sample ( $\geq 200$ )	2
	Clearly identified mixed population	3
	Disorders specific trials (i.e. well defined spinal stenosis and disc herniation, disorder specific, disc herniation or spinal stenosis or post surgery syndrome)	4
7b.	For facet or sacroiliac joint interventions:	
	No specific selection criteria	1
	No diagnostic blocks based on clinical symptomatology	2



## Effectiveness of Therapeutic Facet Joint Interventions

Table 3 (cont.) . *IPM checklist for assessment of nonrandomized or observational studies of IPM techniques utilizing IPM-QRBNR.*

	Selection with single diagnostic blocks	3
	Selection with placebo or dual diagnostic blocks	4
8.	Duration of Pain	
	Less than 3 months	0
	3 to 6 months	1
	> 6 months	2
9.	Previous Treatments	
	Conservative management including drug therapy, exercise therapy, physical therapy, etc.	
	Were not utilized	0
	Were utilized sporadically in some patients	1
	Were utilized in all patients	2
10.	Duration of Follow-up with Appropriate Interventions	
	Less than 3 months or less for epidural or facet joint procedures, etc., and 6 months for intradiscal procedures and implantables	1
	3-6 months for epidural or facet joint procedures, etc., or one year for intradiscal procedures or implantables	2
	6-12 months for epidurals or facet joint procedures, etc., and 2 years or longer for discal procedures and implantables	3
	18 months or longer for epidurals and facet joint procedures, etc., or 5 years or longer for discal procedures and implantables	4
<b>IV. OUTCOMES</b>		
11.	Outcomes Assessment Criteria for Significant Improvement	
	No descriptions of outcomes OR < 20% change in pain rating or functional status	0
	Pain rating with a decrease of 2 or more points or more than 20% reduction OR functional status improvement of more than 20%	1
	Pain rating with decrease of $\geq 2$ points AND $\geq 20\%$ change or functional status improvement of $\geq 20\%$	2
	Pain rating with a decrease of 3 or more points or more than 50% reduction OR functional status improvement with a 50% or 40% reduction in disability score	2
	Significant improvement with pain and function $\geq 50\%$ or 3 points and 40% reduction in disability scores	4
12.	Description of Drop Out Rate	
	No description despite reporting of incomplete data or more than 30% withdrawal	0
	Less than 30% withdrawal in one year in any group	1
	Less than 40% withdrawal at 2 years in any group	2
13.	Similarity of Groups at Baseline for Important Prognostic Indicators	
	No groups or groups dissimilar with significant influence on outcomes	0
	Groups dissimilar without significant influence on outcomes	1
	Groups similar	2
14.	Role of Co-Interventions	
	Dissimilar co-interventions or similar co-interventions in some of the participants	1
	No co-interventions or similar co-interventions in majority of the participants	2
<b>V. ASSIGNMENT</b>		
15.	Method of Assignment of Participants	
	Case report/case series or selective assignment based on outcomes or retrospective evaluation based on clinical criteria	1
	Prospective study with inclusion without specific criteria	2

Table 3 (cont.) . *IPM checklist for assessment of nonrandomized or observational studies of IPM techniques utilizing IPM-QRBNR.*

	Retrospective method with inclusion of all participants or random selection of retrospective data	3
	Prospective, well-defined assignment of methodology and inclusion criteria (quasi randomization, matching, stratification, etc.)	4
<b>VI. CONFLICTS OF INTEREST</b>		
16.	Funding and Sponsorship	
	Trial included industry employees with or without proper disclosure	-3
	Industry employees involved; high levels of funding with remunerations by industry or an organization funded with conflicts	-3
	Industry or organizational funding with reimbursement of expenses with some involvement or no information available	0
	Industry or organization funding of expenses without involvement	1
	Funding by internal resources only	2
	Governmental funding without conflict such as NIH, NHS, AHRQ	3
TOTAL MAXIMUM		48

Source: Manchikanti L, Hirsch JA, Heavner JE, Cohen SP, Benyamin RM, Sehgal N, Falco FJE, Vallejo R, Onyewu O, Zhu J, Kaye AD, Boswell MV, Helm II S, Candido KD, Diwan S, Simopoulos TT, Singh V, Pampati V, Racz GB, Raj PP. Development of an interventional pain management specific instrument for methodologic quality assessment of nonrandomized studies of interventional techniques. *Pain Physician* 2014; 17:E291-E317 (73).

#### **1.4.4 Measurement of Treatment Effect in Data Synthesis (Meta-Analysis)**

If the literature search provided at least 3 randomized trials meeting the inclusion criteria and they are clinically homogenous for each modality and region evaluated, a meta-analysis was performed.

Data were summarized using a meta-analysis when at least 3 trials per type of modality were available that met the inclusion criteria (e.g., intraarticular injections, facet joint nerve blocks, and radiofrequency thermoneurolysis) of clinical and statistical homogeneity.

Qualitative (the direction of a treatment effect) and quantitative (the magnitude of a treatment effect) conclusions were evaluated. A random-effects meta-analysis to pool data was also used. For placebo-controlled trials, the net effect between 2 treatments was utilized. However, for active-controlled trials, the differences between baseline and follow-up period were utilized.

#### **1.5 Outcome of the Studies**

According to reports from trials that studied general chronic pain, a clinically meaningful pain score change is considered to be, at a minimum, a 2-point change on a 0 to 10 scale (or 20 percentage points) (74), chronic musculoskeletal pain (75), and chronic low back pain (71-73,75,76), which have been commonly utilized. Traditional criteria of minimum or meaningful improvements have been criticized as clinically irrelevant (71-73,77-81). Thus, recent descriptions of clinically meaningful improvement considered more robust

outcomes with either pain relief and functional status improvement of 50% (82-97). The following outcomes were considered clinically meaningful or significant: a 3-point or greater change on an 11-point pain scale (0-10), or a 50% pain improvement from baseline and a 40% or greater improvement in functional status.

A trial was judged to be positive if the facet joint intervention was clinically relevant and effective, either with a placebo control or active control. This indicates that the difference in the effect for the primary outcome measure was statistically significant on the conventional 5% level. Negative studies were those where the study treatments showed no difference or there was no improvement from baseline. Outcomes were reported at one, 3, 6, and 12 months. For observational studies, appropriate outcomes were reported with positive or negative results at 3 months, 6 months, and one-year or longer with effectiveness demonstrated when a study was judged to be positive. If a lack of effectiveness was identified in the study, it was judged to be negative.

#### **1.6 Summary Measures**

Summary measures included a 50% or more reduction of pain in at least 50% of the patients, or at least a 3-point decrease in pain scores and a relative risk of adverse events, including side effects.

#### **1.7 Analysis of Evidence**

The analysis of the evidence was performed based on best evidence synthesis developed from modifica-

Table 4. *Qualitative modified approach to grading of evidence.*

<b>Level I</b>	Evidence obtained from multiple relevant high quality randomized controlled trials
<b>Level II</b>	Evidence obtained from at least one relevant high quality randomized controlled trial or multiple relevant moderate or low quality randomized controlled trials
<b>Level III</b>	Evidence obtained from at least one relevant moderate or low quality randomized controlled trial with multiple relevant observational studies or Evidence obtained from at least one relevant high quality nonrandomized trial or observational study with multiple moderate or low quality observational studies
<b>Level IV</b>	Evidence obtained from multiple moderate or low quality relevant observational studies
<b>Level V</b>	Opinion or consensus of large group of clinicians and/or scientists

Source: Manchikanti L, Falco FJE, Benyamin RM, Kaye AD, Boswell MV, Hirsch JA. A modified approach to grading of evidence. *Pain Physician* 2014; 17:E319-E325 (105).

tion of multiple available criteria including those of the United States Preventive Services Task Force (USPSTF) and Cochrane review criteria as illustrated in Table 4 (98-104).

The analysis was conducted utilizing best evidence synthesis using 5 levels of evidence ranging from strong to opinion- or consensus-based (105).

At least 2 of the review authors independently, in an unblinded standardized manner, analyzed the evidence. Any disagreements between reviewers were resolved by a third author and consensus. If there were any conflicts of interest (e.g., authorship), those reviewers were recused from assessment and analysis for the study in question.

## 2.0 RESULTS

Figure 1 shows a flow diagram of the study selection of therapeutic intervention trials and studies.

Based on comprehensive search criteria there were multiple studies considered for inclusion (82-84,106-164).

Multiple randomized trials, duplicates, and all non-randomized trials were excluded. Table 5 is a partial list of excluded trials that did not meet inclusion criteria. Subsequently, 21 randomized trials (82-84,108,110,112-124,128,129) and 5 observational studies were included (152,159-162).

There were 3 trials (82,112,117) that evaluated therapeutic lumbar facet joint nerve blocks, 9 trials (108,113-118,122,123) that evaluated lumbar facet joint radiofrequency neurolysis, and 5 trials (110,111,121,123,124) that evaluated lumbar intraarticular injections that met the inclusion criteria. Even though there were only 3 randomized controlled trials, there were no observational therapeutic facet joint nerve block studies available for inclusion.

There was one trial (128) that evaluated the efficacy of cervical facet joint nerve radiofrequency thermoneurolysis, one trial (83) that evaluated the efficacy of cervical facet joint nerve blocks, and 2 trials (119,129) that evaluated cervical intraarticular injections that met inclusion criteria. Thus, 3 observational studies assessing radiofrequency neurotomy of the cervical spine (159-161) were included in the assessment of radiofrequency neurotomy. In addition, one prospective evaluation of cervical facet joint nerve blocks was also included (152).


There was one trial (84) that evaluated therapeutic thoracic facet joint nerve blocks and one trial (120) that evaluated thoracic facet joint radiofrequency neurolysis that met the inclusion criteria. Thus, one study of thoracic facet joint nerve blocks (162) was included. There were no other studies meeting inclusion criteria.

### 2.1 Methodological Quality Assessment

A methodological quality assessment of the randomized controlled trials meeting inclusion criteria was carried out utilizing Cochrane review criteria and IPM-QRB criteria for randomized trials as shown in Tables 6 and 7 and IPM – QRBNR for nonrandomized studies as shown in Table 8.

Utilizing Cochrane review criteria, studies meeting the inclusion criteria with at least a score of 8 of 12 were considered high quality and a score of 4 to 7 were considered moderate quality. A score lower than 4 was considered low quality and those studies were excluded.

Based on IPM-QRB criteria for randomized trials and IPM-QRBNR for observational studies, the trials meeting the inclusion criteria with scores of less than 16 were considered low quality and were excluded, manuscripts with scores of 16 to 31 were considered moderate quality, and scores of 32 to 48 or higher were considered as high quality trials.



## 2.2 Meta-Analysis

If there were more than 2 trials meeting clinical homogeneity criteria, they were further assessed for homogeneity, and a meta-analysis was performed. There were 9 trials assessing lumbar radiofrequency neurotomy (108,113-118,122,123), 5 trials assessing lumbar

intraarticular injection therapy (110,111,121,123,124), and 3 trials assessing lumbar facet joint nerve blocks (82,112,117). However, all modalities in the cervical spine and the thoracic spine had 2 or fewer. An assessment of clinical and methodological homogeneity

## Effectiveness of Therapeutic Facet Joint Interventions

Table 5. List of excluded randomized trials with brief explanation.

Study	Condition Studied	Number of Patients	Reason for Exclusion	
			Followup Period	Other Reason(s)
<b>LUMBAR FACET JOINT INTERVENTIONS</b>				
Lilius et al (131) Randomized	Chronic low back pain	109	3 months	Study with short-term follow-up along with lack of diagnostic blocks and comparison of intraarticular or extraarticular injections with a large volume of injection. At best, this study may be appropriate for a diagnostic study with a single block.
Marks et al (132) Randomized	Chronic low back pain	86	3 months	The authors compared facet joint nerve blocks and intraarticular injections with high volume injections with very short-term follow-up in a randomized trial as diagnostic blocks.
Nash (133) Randomized	Chronic low back pain	67	3 months	The authors compared the effectiveness of intraarticular injections with medial branch blocks on a short-term basis with no controlled local anesthetic blocks, and with lack of long-term follow-up and outcomes
Leclaire et al (134) Randomized, Placebo-Controlled	Chronic low back pain	70	12 weeks	Relatively small study; however, technique and the diagnostic evaluation with intraarticular injections were inappropriate. The authors have admitted that the results might not be applicable in clinical practice.
Gallagher et al (135) Randomized	Chronic low back pain	41	One month and 6 months	Authors evaluated 60 patients with a single block and randomized them into 2 groups with 41 patients testing positive. The study showed improvement at one month and 6 months; however, the inclusion criteria, the technical considerations, and statistical analysis were considered as flawed.
Kroll et al (136) Randomized	Acute low back pain	50	3 months	Conventional and pulsed radiofrequency neurotomy were studied in acute low back pain.
Kader et al (140) Randomized	Chronic nonspecific low back pain with or without leg pain	63	10 weeks	Patients were randomized into 3 groups with back education and standard physiotherapy for 10 weeks, back education and gym ball exercise for 10 weeks, or perifacet injection into the lumbar multifidus muscle with methylprednisolone. Since there was no facet joint injection, the study failed to meet the criteria for inclusion.
Wen et al (126) Randomized	Low back pain	20	6 weeks	Twenty with 10 patients in each group receiving facet joint blocks either blindly or guided by ultrasound; however, the needle tip was confirmed by CT in both groups. Small sample size with short-term follow-up of 6 weeks utilizing either a blind technique or ultrasound-guided lumbar facet joint injections.
<b>CERVICAL FACET JOINT INTERVENTIONS</b>				
Obernauer et al (125) Randomized	Subacute chronic facet-joint-associated neck pain of the middle or lower cervical spine	40	4 weeks	Injection of local anesthetic and steroids into cervical facet joints either under CT scanning or ultrasonography. The short-term follow-up in patients with subacute pain was done without diagnostic blocks.
Slappendel et al (148) Randomized	Cervicobrachialgia	61	3 months	The authors evaluated the efficacy of radiofrequency lesioning of the cervical dorsal root ganglion.
Haspeslagh et al (149) Randomized	Cervicogenic headache	30	48 weeks	In this study, 30 patients with cervicogenic headache were evaluated. This study was problematic, not only in the diagnosis but also in the application of technique.  The authors claim that they developed a sequence of various cervical radiofrequency neurotomies that proved successful in a prospective pilot trial with 15 chronic headache patients. Their diagnosis was not established by controlled diagnostic blocks; and the treatments targeted toward different structures: cervical facet joints and dorsal root ganglia compared to occipital nerves.

Table 6. Methodological quality assessment of randomized trials of lumbar, cervical and thoracic facet joint interventions utilizing Cochrane review criteria.

	Manchikanti et al (82)	Carette et al (110)	Fuchs et al (111)	Nath et al (113)	van Wijk et al (114)	van Kleef et al (115)	Tekin et al (116)	Givolek et al (117)	Dobrogowski et al (118)	Cohen et al (108)	Barnsley et al (119)
Randomization adequate	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Concealed treatment allocation	Y	Y	N	Y	Y	Y	Y	Y	U	N	Y
Patient blinded	Y	Y	Y	Y	Y	Y	Y	N	Y	N	Y
Care provider blinded	Y	Y	N	Y	Y	Y	Y	N	Y	U	Y
Outcome assessor blinded	N	Y	Y	Y	Y	Y	Y	U	U	U	Y
Drop-out rate described	Y	Y	N	Y	Y	Y	Y	Y	Y	Y	Y
All randomized participants analyzed in the group	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Reports of the study free of suggestion of selective outcome reporting	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Groups similar at baseline regarding most important prognostic indicators	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Co-intervention avoided or similar in all groups	Y	N	N	Y	Y	Y	Y	Y	Y	Y	Y
Compliance acceptable in all groups	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Time of outcome assessment in all groups similar	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<b>SCORE</b>	11/12	11/12	8/12	12/12	12/12	12/12	12/12	9/12	10/12	8/12	12/12
	Manchikanti et al (83)	Lord et al (128)	Park & Kim (129)	Manchikanti et al (84)	Manchikanti et al (120)	Ribeiro et al (121)	Moon et al (122)	Lakemeier et al (123)	Yun et al (124)	Manchikanti et al (112)	
Randomization adequate	Y	Y	N	Y	Y	Y	Y	Y	Y	N	
Concealed treatment allocation	Y	Y	N	Y	Y	Y	Y	Y	Y		
Patient blinded	Y	Y	N	Y	Y	Y	Y	Y	N	Y	
Care provider blinded	Y	Y	N	Y	N	N	Y	N	N	Y	
Outcome assessor blinded	N	Y	N	N	N	N	Y	N	N	N	
Drop-out rate described	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	
All randomized participants analyzed in the group	Y	Y	Y	Y	Y	Y	N	Y	Y	N	
Reports of the study free of suggestion of selective outcome reporting	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	
Groups similar at baseline regarding most important prognostic indicators	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	
Co-intervention avoided or similar in all groups	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	
Compliance acceptable in all groups	Y	Y	N	Y	Y	Y	N	N	Y	N	
Time of outcome assessment in all groups similar	Y	N	Y	Y	Y	Y	N	Y	Y	Y	
<b>SCORE</b>	11/12	11/12	6/12	10/12	10/12	10/12	9/12	9/12	9/12	6/12	

Source: Furlan AD, Pennick V, Bombardier C, van Tulder M; Editorial Board, Cochrane Back Review Group. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine (Phila Pa 1976)* 2009; 34:1929-1941 (70).

Effectiveness of Therapeutic Facet Joint Interventions

Table 7. Methodologic quality assessment of randomized trials of lumbar, cervical and thoracic facet joint interventions utilizing IPM – QRB criteria.

		Manchikanti et al (82)	Carette et al (110)	Fuchs et al (111)	Nath et al (113)	van Wijk et al (114)	van Kleef et al (115)	Tekin et al (116)	Givelek et al (117)	Dobrogowski et al (118)	Cohen et al (108)
I.	Trial design and guidance reporting										
1.	Consort or spirit	3	3	3	3	2	2	2	2	2	3
II.	Design factors										
2.	Type and design of trial	2	3	2	3	3	3	3	2	2	2
3.	Setting/physician	2	1	1	3	3	3	2	2	2	2
4.	Imaging	3	3	2	3	3	3	3	3	2	3
5.	Sample size	3	2	2	1	1	1	1	1	1	3
6.	Statistical methodology	1	1	1	1	1	1	1	1	1	1
III.	Patient factors										
7.	Inclusiveness of population										
	• For facet or sacroiliac joint interventions:	2	1	0	2	1	1	1	0	1	1
8.	Duration of pain	2	2	1	2	2	3	2	0	2	0
9.	Previous treatments	2	0	0	0	0	0	1	2	0	0
10.	Duration of follow-up with appropriate interventions	3	2	1	2	2	2	2	2	1	0
IV.	Outcomes										
11.	Outcomes assessment criteria for significant improvement	4	4	2	4	4	2	2	2	2	0
12.	Analysis of all randomized participants in the groups	2	2	2	2	2	2	2	2	2	2
13.	Description of drop out rate	2	1	2	2	2	2	2	2	2	2
14.	Similarity of groups at baseline for important prognostic indicators	2	2	2	2	2	2	2	2	2	2
15.	Role of co-interventions	1	0	0	1	1	1	1	1	1	1
V.	Randomization										
16.	Method of randomization	2	2	2	2	2	2	2	2	2	2
VI.	Allocation concealment										
17.	Concealed treatment allocation	2	2	1	2	2	2	2	2	2	0
VII.	Blinding										
18.	Patient blinding	1	1	0	1	1	1	1	0	1	0
19.	Care provider blinding	1	1	0	1	1	1	1	0	1	0
20.	Outcome assessor blinding	0	1	0	0	1	0	0	0	0	0
VIII.	Conflicts of interest										
21.	Funding and sponsorship	2	3	1	2	0	3	2	0	0	2
22.	Conflicts of interest	3	3	1	3	0	3	2	0	0	2
TOTAL		45	40	26	42	36	40	37	28	29	28

Source: Manchikanti L, Hirsch JA, Cohen SP, Heavner JE, Falco FJE, Diwan S, Boswell MV, Candido KD, Onyewu O, Zhu J, Sehgal N, Kaye AD, Benjamin RM, Helm II S, Singh V, Datta S, Abdi S, Christo PJ, Hameed H, Hameed M, Vallejo R, Pampati V, Racz GB, Raj PP. Assessment of methodologic quality of randomized trials of interventional techniques: Development of an interventional pain management specific instrument. Pain Physician 2014; 17:E263-E290 (72).

Table 7 (cont.). Methodologic quality assessment of randomized trials of lumbar, cervical and thoracic facet joint interventions utilizing IPM – QRB criteria.

	Barnsley et al (119)	Manchikanti et al (83)	Lord et al (128)	Park & Kim (129)	Manchikanti et al (84)	Joo et al (120)	Ribeiro et al (121)	Moon et al (122)	Lakemeier et al (123)	Yun et al (124)	Manchikanti et al (112)	
<b>I. Trial design and guidance reporting</b>												
1.	Consort or spirit	2	3	3	2	3	2	2	2	2	2	2
<b>II. Design factors</b>												
2.	Type and design of trial	2	2	3	2	2	2	2	2	2	2	2
3.	Setting/physician	2	2	2	2	2	2	2	2	1	1	3
4.	Imaging	3	3	3	3	3	3	3	3	3	3	3
5.	Sample size	1	3	1	3	3	1	2	2	2	2	2
6.	Statistical methodology	1	1	1	1	1	1	1	1	1	1	1
<b>III. Patient factors</b>												
7.	Inclusiveness of population											
	• For facet or sacroiliac joint interventions:	2	2	2	2	2	2	0	2	1	0	2
8.	Duration of pain	2	2	2	2	2	2	0	2	2	0	2
9.	Previous treatments	0	2	2	2	2	2	0	2	2	1	2
10.	Duration of follow-up with appropriate interventions	1	3	2	2	3	2	1	1	2	1	3
<b>IV. Outcomes</b>												
11.	Outcomes assessment criteria for significant improvement	2	4	4	2	4	2	2	2	2	1	2
12.	Analysis of all randomized participants in the groups	2	2	2	2	2	2	2	0	2	2	0
13.	Description of drop out rate	1	2	2	2	2	2	2	2	2	2	0
14.	Similarity of groups at baseline for important prognostic indicators	2	2	2	2	2	2	2	2	2	2	2
15.	Role of co-interventions	0	1	1	0	1	1	1	1	1	1	1
<b>V. Randomization</b>												
16.	Method of randomization	2	2	2	2	2	2	2	2	2	2	0
<b>VI. Allocation concealment</b>												
17.	Concealed treatment allocation	2	2	2	0	2	2	2	2	2	2	0
<b>VII. Blinding</b>												
18.	Patient blinding	1	1	1	0	1	1	1	1	1	0	1
19.	Care provider blinding	1	1	1	0	1	0	0	1	0	0	1



## Effectiveness of Therapeutic Facet Joint Interventions

*Table 7 (cont.). Methodologic quality assessment of randomized trials of lumbar, cervical and thoracic facet joint interventions utilizing IPM – QRB criteria.*

		Barnsley et al (119)	Manchikanti et al (83)	Lord et al (128)	Park & Kim (129)	Manchikanti et al (84)	Joo et al (120)	Ribeiro et al (121)	Moon et al (122)	Lakemeier et al (123)	Yun et al (124)	Manchikanti et al (112)
20.	Outcome assessor blinding	1	0	1	0	0	0	0	1	0	0	0
<b>VIII. Conflicts of interest</b>												
21.	Funding and sponsorship	3	2	3	2	2	2	2	2	2	1	2
22.	Conflicts of interest	3	3	3	2	3	3	3	3	3	0	3
<b>TOTAL</b>		<b>36</b>	<b>45</b>	<b>45</b>	<b>35</b>	<b>45</b>	<b>38</b>	<b>32</b>	<b>38</b>	<b>37</b>	<b>26</b>	<b>34</b>

Source: Manchikanti L, Hirsch JA, Cohen SP, Heavner JE, Falco FJE, Diwan S, Boswell MV, Candido KD, Onyewu O, Zhu J, Sehgal N, Kaye AD, Benjamin RM, Helm II S, Singh V, Datta S, Abdi S, Christo PJ, Hameed H, Hameed M, Vallejo R, Pampati V, Racz GB, Raj PP. Assessment of methodologic quality of randomized trials of interventional techniques: Development of an interventional pain management specific instrument. *Pain Physician* 2014; 17:E263-E290 (72).

*Table 8. IPM checklist for assessment of nonrandomized or observational studies of lumbar, cervical and thoracic facet joint interventions of IPM techniques utilizing IPM-QRBNR.*

		Sapir & Gorup (159)	MacVicar et al (160)	Speldewinde GC (161)	Manchikanti et al (152)	Manchikanti et al (162)
<b>I.</b>	<b>Study design and guidance reporting</b>					
1.	Strobe or trend guidance	3	3	3	3	3
<b>II.</b>	<b>Design factors</b>					
2.	Study design and type	4	4	4	4	4
3.	Setting/physician	2	2	2	2	2
4.	Imaging	3	3	3	3	3
5.	Sample size	2	1	1	1	1
6.	Statistical methodology	2	2	2	2	2
<b>III.</b>	<b>Patient factors</b>					
7.	Inclusiveness of population					
	• For facet or sacroiliac joint interventions:	4	4	4	4	4
8.	Duration of pain	2	2	2	2	2
9.	Previous treatments	2	2	2	2	2
10.	Duration of follow-up with appropriate interventions	3	3	3	3	3
<b>IV.</b>	<b>Outcomes</b>					
11.	Outcomes assessment criteria for significant improvement	2	4	4	4	4
12.	Description of drop out rate	1	1	1	1	1
13.	Similarity of groups at baseline for important prognostic indicators	2	0	0	0	0
14.	Role of co-interventions	2	2	2	2	2
<b>V.</b>	<b>Assignment</b>					
15.	Method of assignment of participants	4	4	4	2	2
<b>VI.</b>	<b>Conflicts of interest</b>					
16.	Funding and sponsorship	2	1	2	2	2
<b>TOTAL</b>		<b>40</b>	<b>38</b>	<b>39</b>	<b>37</b>	<b>37</b>

Source: Manchikanti L, Hirsch JA, Heavner JE, Cohen SP, Benjamin RM, Sehgal N, Falco FJE, Vallejo R, Onyewu O, Zhu J, Kaye AD, Boswell MV, Helm II S, Candido KD, Diwan S, Simopoulos TT, Singh V, Pampati V, Racz GB, Raj PP. Development of an interventional pain management specific instrument for methodologic quality assessment of nonrandomized studies of interventional techniques. *Pain Physician* 2014; 17:E291-E317 (73).

among various studies of the 11 trials of radiofrequency neurotomy was performed. None of the 3 trials were homogeneous either for selection criteria, outcome measures, or design of the trial. Among the 5 lumbar intraarticular injections and 3 facet joint nerve blocks there was no clinical homogeneity in any of the 2 or more trials. Thus, a meta-analysis was not feasible; consequently it was not performed.

### 2.3 Study Characteristics

Tables 9 and 10 illustrate the study characteristics of the included studies for randomized trials and observational studies evaluating facet joint interventions.

### 2.4 Analysis of Evidence

The evidence was synthesized based on the modality of treatment for each region. Tables 11 and 12 illustrate the results of therapeutic studies.

A total of 21 randomized trials met inclusion criteria with 9 trials (108,113-118,122,123) evaluating lumbar radiofrequency neurotomy, one trial evaluating cervical radiofrequency neurotomy (128), and one trial evaluating thoracic radiofrequency neurotomy (120); with 3 trials (82,112,117) evaluating therapeutic lumbar facet joint nerve blocks, one trial evaluating therapeutic cervical facet joint nerve blocks (83), and one trial evaluating therapeutic thoracic facet joint nerve blocks (84); and 5 trials evaluating lumbar intraarticular injections (110,111,121,123,124) and 2 trials evaluating cervical intraarticular injections (119,129). In addition, one additional therapeutic thoracic facet joint nerve blocks study (162), one additional therapeutic cervical facet joint nerve block study (152), and 3 additional cervical radiofrequency neurotomy studies (159-161) were also included.

The evidence for radiofrequency neurotomy in the lumbar, cervical, and thoracic spines is variable. The evidence is Level I for short-term effectiveness of radiofrequency neurotomy of less than 6 months and Level II for long-term relief of longer than 6 months based on 8 moderate to high quality randomized controlled trials of radiofrequency neurotomy (108,113,115,116,117,118,122,123) showing short-term effectiveness of radiofrequency neurotomy, lack of response shown in one moderate to high quality trial of radiofrequency neurotomy (114), long-term relief based on 3 high quality randomized controlled trials showing effectiveness (115-117), lack of effectiveness demonstrated in one trial (114). In the cervical spine, the evidence is Level II based on one high quality

randomized controlled trial (128) for short-term and long-term effectiveness; whereas, in the thoracic spine the evidence is Level III based on one randomized, double-blind, active control trial (120) for short-term and long-term effectiveness.

The evidence for therapeutic facet joint nerve blocks is Level II in the cervical, thoracic, and lumbar regions for short-term and long-term improvement based on 2 high quality randomized controlled trials (82,117) and one moderate (6/12 Cochrane criteria) to high quality (34/48 IPM-QRB criteria) randomized controlled trial (112) of facet joint nerve blocks in the lumbar spine with long-term follow-up (82,117). In the cervical spine, the evidence is Level II based on one high quality randomized controlled trial (83) and the evidence is Level II in the thoracic spine based on one high quality randomized controlled trial (84).

The evidence for intraarticular facet joint injections is variable between the lumbar and cervical spines. There is no evidence available for thoracic intraarticular injections. The evidence for lumbar intraarticular injections of steroids is Level III, based on 3 high quality randomized controlled trials (121,123,124) showing effectiveness with short-term follow-up of less than 6 months and 2 moderate to high quality randomized controlled trials (110,111) showing a lack of effectiveness with a follow-up shorter than 6 months for short-term pain relief. The evidence for cervical intraarticular injections is Level IV based on one high quality randomized controlled trial (119) showing a lack of effectiveness and one moderate quality randomized controlled trial (129) demonstrating indeterminate results.

## 3.0 DISCUSSION

The systematic review of randomized trials of efficacy of the spinal facet joint interventions in the lumbar, cervical and thoracic regions, with intraarticular injections, facet joint nerve blocks, and radiofrequency neurotomy, revealed variable results. A total of 20 randomized trials were assessed with moderate to high methodologic quality criteria.

Based on the available evidence, there is Level II evidence for lumbar radiofrequency neurotomy, Level II evidence for cervical radiofrequency neurotomy, and Level III evidence for thoracic radiofrequency neurotomy for long-term effectiveness. The evidence is Level II for lumbar, cervical, and thoracic facet joint nerve blocks for long-term effectiveness. For intraarticular injections, the evidence is Level III for lumbar intraarticular injections and Level IV for cervical intraarticular injections;

Table 9. Study characteristics of randomized controlled trials and observational studies assessing lumbar radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>RADIOFREQUENCY NEUROTOMY</b>									
Civelek et al, 2012 (117) Randomized, active-control trial Quality Scores: Cochrane = 9/12 IPM-QRB = 28/48	100 patients with chronic low back pain with failed conservative therapy and strict selection criteria; however, without diagnostic blocks.	Facet joint nerve block with local anesthetic and steroids in 50 patients.	Conventional radiofrequency neurotomy at 80°C for 120 seconds in combination with high dose local anesthetic and steroids, in 50 patients.	Visual Numeric Pain Scale, North American Spine Society patient satisfaction questionnaire, Euro-Qol in 5 dimensions and ≥ 50% relief	One month, 6 months, 12 months	At one year, 90% of patients in the radiofrequency group and 69% of the patients in the facet joint nerve block group showed significant improvement compared to 92% and 75% at 6-month follow-up.	Randomized relatively large number of patients with 50 in each group.	No diagnostic blocks were performed. High dose steroids and local anesthetics were utilized in both groups.	Efficacy was shown even without diagnostic blocks, both for facet joint nerve blocks and radiofrequency neurotomy.
Cohen et al, 2010 (108) Randomized, double-blind, active control trial Quality Scores: Cochrane = 8/12 IPM-QRB = 28/48	151 chronic low back pain 51 patients with no diagnostic block 50 patients a single diagnostic block 50 patients in double diagnostic block.	Radiofrequency neurotomy in patients without diagnostic blocks.	Conventional radiofrequency neurotomy at 80°C for 90 seconds in all patients; however, in 2 groups with either a single block paradigm or a double block paradigm testing for positive results.	Greater than 50% pain relief coupled with a positive global perceived effect persisting for 3 months.	3 months	Denervation success rates in Groups 0, 1, and 2 were 33%, 39%, and 64% respectively.	Multicenter, randomized controlled trial with or without diagnostic blocks	Authors misinterpreted cost-effectiveness without consideration of many factors reported.	Results showed efficacy when double diagnostic blocks were utilized.
Nath et al, 2008 (113) Randomized, double-blind, sham control trial Quality Scores: Cochrane = 12/12 IPM-QRB = 42/48	40 patients with chronic low back pain for at least 2 years with 80% relief of low back pain after controlled medial branch blocks. The patients were randomized into an active and a control group.	Sham control with placement of the needles with injection of local anesthetic without radiofrequency neurotomy.	The 20 patients in the active group received conventional lumbar facet joint radiofrequency neurolysis at 85°C for 60 seconds. The 20 patients in the control group received sham treatment without radiofrequency neurolysis of the lumbar facet joints.	Numeric Rating Scale, global functional improvement, reduced opioid intake, employment status.	6 months	Significant reduction not only in back, and leg pain; functional improvement; opioid reduction; and employment status in the active group compared to the control group.	Randomized, double-blind trial after the diagnosis of facet joint pain with triple diagnostic blocks	Short-term follow-up with small number of patients	Efficacy of radiofrequency neurotomy was shown compared to local anesthetic injection and sham lesioning.
Tekin et al, 2007 (116) Randomized, active and sham, double-blind controlled trial Quality Scores: Cochrane = 12/12 IPM-QRB = 37/48	60 patients with chronic low back pain randomized into 3 groups with 20 patients in each group. Single diagnostic block of facet joint nerves with 0.3 mL of lidocaine 2% with 50% or greater relief.	Sham control with local anesthetic injection	Either pulsed radiofrequency (42°C for 4 minutes) or conventional radiofrequency neurotomy (80°C for 90 seconds) in 20 patients in each group.	Visual analog scale and Oswestry Disability Index	3, 6, and 12 months	Visual analog scale and Oswestry Disability Index scores decreased in all groups from 3 procedural levels. Decrease in pain scores was maintained in the conventional radiofrequency group at 6 months and one year. However, in pulsed radiofrequency group, the improvement was significant only at 6 months, but not one year.	Randomized, double-blind, controlled trial comparing control, pulsed radiofrequency, and conventional radiofrequency neurotomy. Authors also utilized a parallel needle placement approach	Small sample size with a single block and 50% relief as inclusion criteria. Authors did not report significant improvement percentages.	Efficacy with conventional radiofrequency neurotomy up to one year, whereas efficacy with local anesthetic block with sham control radiofrequency neurotomy and pulsed radiofrequency neurotomy at 6 months only.

Table 9 (cont.). Study characteristics of randomized controlled trials and observational studies assessing lumbar radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Study Characteristic Methodological Quality Scoring	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
van Wijk et al, 2005 (114)  Randomized, double-blind, sham control trial  Quality Scores: Cochrane = 12/12 IPM-QRB = 36/48	81 patients with chronic low back pain were evaluated with radiofrequency neurotomy with 41 patients in the control group with at least 50% relief for 30 minutes with a single block with intraarticular injection of 0.5 mL lidocaine 2%.	Sham lesion procedure after local anesthetic injection	40 patients received conventional radiofrequency lesioning at 80°C for 60 seconds and 41 patients received sham lesioning.	Pain relief, physical activities, analgesic intake, global perceived effect. Short-form-36, quality of life measures	3 months	Global perceived effect improved after radiofrequency facet joint denervation. The visual analog scale in both groups improved. The combined outcome measures showed no difference between radiofrequency facet joint denervation (27.5% vs. 29.3% success rate).	Double-blind, sham control, randomized trial	Poor selection with a single diagnostic block of 50% pain reduction even though 17.5% of the patients were tested positive. Further, authors described that the needle was positioned parallel; however, the radiographic figures illustrate the needle was being positioned perpendicularly rather than parallel to the nerve.	Lack of efficacy with methodologic deficiencies and a short-term follow-up.	
Dobrogowski et al, 2005 (118)  Randomized, active control trial  Quality Scores: Cochrane = 10/12 IPM-QRB = 29/48	45 consecutive patients with chronic low back pain judged to be positive with significant relief with 2 controlled diagnostic blocks	Injection of saline in patients after conventional radiofrequency (85° for 60 seconds) neurotomy to evaluate postoperative pain	Conventional radiofrequency neurotomy at 85°C for 60 seconds, followed by injection of either methylprednisolone or pentoxifylline	Visual analog scale, minimum of 50% reduction of pain intensity, patient satisfaction score	One, 3, 6, and 12 months	Greater than 50% reduction of pain intensity was observed in 66% of the patients 12 months later. There was no difference in the long-term outcomes.	Randomized, active control trial	Very small study with highly defined inclusion criteria evaluating effectiveness of radiofrequency neurotomy and postoperative pain.	Radiofrequency neurotomy effective with or without steroid injection after neurolysis.	
Van Kleef et al, 1999 (115)  Randomized, double-blind, sham control trial  Quality Scores: Cochrane = 12/12 IPM-QRB = 40/48	31 patients with a history of at least one year of chronic low back pain randomly assigned to one of 2 treatment groups. Single diagnostic block with 50% relief.	Sham control of radiofrequency after local anesthetic injection in 16 patients	The 15 patients in the conventional radiofrequency treatment group received an 80° C radiofrequency lesion for 60 seconds.	Visual analog scale, pain scores, global perceived effect, Oswestry Disability Index	3, 6, and 12 months	After 3, 6, and 12 months, the number of successes in the lesion and sham groups was 9 of 15 (60%) and 4 of 16 (25%), 7 of 15 (47%) and 3 of 16 (19%), and 7 of 15 (47%) and 2 of 16 (13%) respectively. There was a statistically significant difference.	Double-blind, randomized, sham controlled trial	A single block with a small sample with inclusion criteria of 50% pain relief to enter the study. The study has been criticized that electrodes were placed at an angle to the target nerve, instead of parallel.	Efficacy shown in a small sample with a single diagnostic block	

Table 9 (cont.). Study characteristics of randomized controlled trials and observational studies assessing lumbar radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<p>Moon et al, 2013 (122)</p> <p>Prospective, randomized, active control, comparative analysis</p> <p>Quality Scores: Cochrane = 9/12 IPM-QRB = 38/48</p>	<p>82 patients were included with low back pain with 41 patients in each group either with a parallel placement of the needle or perpendicular placement of the needle.</p> <p>Concordant pain relief of &gt;50% after a comparative local anesthetic block.</p>	<p>An active control trial with needle placement with perpendicular approach.</p>	<p>41 patients in each group were treated with radiofrequency (80°C for 90 seconds) after appropriate diagnosis of facet joint pain with dual diagnostic blocks with 50% relief as the criterion standard. The needle was positioned either utilizing a discal or perpendicular approach or utilizing a tunnel vision approach with parallel placement of the needle.</p>	<p>Numeric Rating Scale, Oswestry Disability Index</p>	<p>One month and 6 months</p>	<p>Patients in both groups showed a statistically significant reduction in NRS and Oswestry disability index scores from baseline to that of the scores at one and 6 months (all <math>P &lt; 0.0001</math>, Bonferroni corrected).</p>	<p>Randomized, double-blind, controlled trial. The major strength is that authors have proven that parallel approach may not be the best as has been described. Diagnosis of facet joint pain by dual blocks.</p>	<p>Active controlled trial without placebo group. Short-term follow-up.</p>	<p>Positive results in an active controlled trial, in a relatively short-term follow-up of 6 months, with positioning of the needle either with distal approach (perpendicular placement or tunnel vision) with parallel placement of the needle with some superiority with perpendicular approach. This trial abates any criticism of needle positioning one way or the other and the traditional needle positioning appears to be superior to parallel needle placement.</p>
<p>Lakemeier et al, 2013 (123)</p> <p>Randomized, double-blind, active controlled trial</p> <p>Quality Scores: Cochrane = 9/12 IPM-QRB = 37/48</p>	<p>56 patients were randomized into 2 groups with 29 patients receiving intraarticular steroid injections and 27 patients receiving radiofrequency denervation after the diagnosis was made with intraarticular injection of local anesthetic with a single block with pain reduction of at least 50%.</p>	<p>Intraarticular injection of local anesthetic and steroid</p>	<p>Radiofrequency neurotomy for 90 seconds at 80°C</p>	<p>Roland-Morris questionnaire, visual analog scale, Oswestry Disability Index, analgesic intake</p>	<p>6 months</p>	<p>Pain relief and functional improvement were observed in both groups. There were no significant differences between the 2 groups for pain relief and functional status improvement.</p>	<p>Lack of placebo group. Relatively short-term follow-up.</p>	<p>Randomized, double-blind trial with single diagnostic block with intraarticular injection</p>	<p>Both groups showed improvement. Effectiveness at 6 months in both groups with intraarticular injection or radiofrequency neurotomy.</p>

Table 9 (cont.). Study characteristics of randomized controlled trials and observational studies assessing lumbar radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>LUMBAR FACET JOINT NERVE BLOCKS</b>									
Givalek et al, 2012 (117)	100 patients with chronic low back pain with failed conservative therapy and strict selection criteria; however, without diagnostic blocks.	Blocks of facet joint nerves with local anesthetic and steroids.	Conventional radiofrequency neurotomy at 80°C for 120 seconds in combination with high dose local anesthetic and steroids.	Visual Numeric Pain Scale, North American Spine Society patient satisfaction questionnaire, Euro-Qol in 5 dimensions and ≥ 50% relief	One month, 6 months, 12 months	At the end of one year, 90% of patients in the radiofrequency group and 69% of the patients in the facet joint nerve block group showed significant improvement vs. 92% and 75% at 6-month follow-up.	Randomized active-control trial with relatively large number of patients with 50 in each group.	No diagnostic blocks were performed. High dose steroids and local anesthetics were provided in both groups.	Results showed efficacy even without diagnostic blocks, both for facet joint nerve blocks and radiofrequency neurotomy.
Manchikanti et al, 2010 (82)	120 patients with chronic low back pain of facet joint origin treated with therapeutic lumbar facet joint nerve blocks.	Local anesthetic only	Total of 120 patients with 60 patients in each group with local anesthetic alone or local anesthetic and steroids. Both groups were also divided into 2 categories each with the addition of Sarapin.	Numeric Rating Scale, Oswestry Disability Index, employment status, and opioid intake.	3, 6, 12, 18, and 24 months	Significant pain relief was shown in 85% in local anesthetic group and 90% in local anesthetic with steroids group at the end of the 2 year study period in both groups, with an average of 5-6 total treatments.	Randomized trial with relatively large proportion of patients with 2-year follow-up, with inclusion of patients diagnosed with controlled diagnostic blocks	Lack of placebo group	Effectiveness demonstrated with facet joint nerve blocks with local anesthetic with or without steroids.
Manchikanti et al, 2001 (112)	73 patients with chronic low back pain diagnosed with dual diagnostic blocks were selected	Active control with local anesthetic and Sarapin	Facet joint nerve blocks with local anesthetic and Sarapin or with local anesthetic, Sarapin, and methylprednisolone	Numeric pain rating scale, functional status, opioid intake, employment status	2 ½ years	Results showed significant improvement in patients in both groups. Significant relief was seen in 100% of patients up to 3 months, 75% in Group I and 80% in Group II at 6 months and 12 months.	Moderate quality randomized trial with long-term follow-up	Appropriate randomization and allocation concealment were lacking	Positive results with cost utility
<b>LUMBAR INTRAARTICULAR INJECTIONS</b>									
Carette et al, 1991 (110)	Patients with chronic low back pain who reported immediate relief of their pain after injection of local anesthetic into the facet joints. Single diagnostic blocks with 50% relief were randomly assigned to receive injections under fluoroscopic guidance.	Intraarticular injection of isotonic saline	Injection of either sodium chloride or methylprednisolone into the facet joints (49 for isotonic saline and 48 for sodium chloride). Only one injection was provided.	Visual Analog Scale, McGill Pain Questionnaire, mean sickness impact profile.	One, 3, and 6 months	After one month, 42% of the patients in the methylprednisolone group and 33% in the sodium chloride group reported marked or very marked improvement. At the 6 month evaluation, 46% in the methylprednisolone group and 15% in the placebo group showed sustained relief. Revised statistics showed 22% improvement in active group and 10% in control group.	Well-performed randomized, double-blind controlled trial	Only single block was applied and patients were treated with steroids without local anesthetic with only one treatment and expected 6 months of relief.	The authors concluded that results were negative in an active-control trial with injection of either sodium chloride solution or steroid into the facet joints after diagnosis with a single block.

Table 9 (cont.). Study characteristics of randomized controlled trials and observational studies assessing lumbar radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
Fuchs et al, 2005 (111) Randomized, double-blind, active-control trial Quality Scores: Cochrane = 8/12 IPM-QRB = 26/48	60 patients with chronic low back pain were included with patients randomly assigned into 2 groups. No diagnostic blocks	Active-control study with no control group	Intraarticular injection of hyaluronic acid versus glucocorticoid injection.	Visual Analog Scale, Rowland-Morris Questionnaire, Oswestry Disability Index, low back outcomes score, Short Form-36	3 months and 6 months	Patients reported lasting pain relief, better function, and improved quality of life with both treatments.	Randomized, active-control, double-blind study	Relatively small sample of patients with 6 month follow-up without a placebo group, without diagnostic blocks.	Undetermined (clinically inapplicable) results with high number of injections during a 6-month period.
Ribeiro et al, 2013 (121) Randomized, double-blind, active control Quality Scores: Cochrane = 10/12 IPM-QRB = 32/48	60 patients with a diagnosis of facet joint syndrome randomized into experimental and control groups.	Triamcinolone acetamide intramuscular injection of 6 lumbar paravertebral points	Intraarticular injection of 6 lumbar facet joints with triamcinolone hexacetone	Pain visual analog scale, pain visual analog scale during extension of the spine, Likert scale, improvement percentage, Roland-Morris, 36-Item Short Form Health Survey, and accountability of medications taken.	One, 4, 12, and 24 weeks	Improvement "percentage" analysis at each time point, showed significant differences between the groups at week 7 and week 12. Improvement percentage was > 50% at all times in the experimental group with intraarticular steroids; however, significant difference was noted at 24 weeks only.	Randomized, double-blind controlled trial	Diagnostic blocks were not employed, thus, many patients without facet joint pain may have been included in this trial.	Overall intraarticular steroids showed positive effective for 24 weeks compared to intramuscular steroids provided in a double-blind manner.
Lakemeier et al, 2013 (123) Randomized, double-blind, active controlled trial Quality Scores: Cochrane = 9/12 IPM-QRB = 37/48	56 patients were randomized into 2 groups receiving intraarticular steroid injections or radiofrequency denervation after the diagnosis was made with intraarticular injection of local anesthetic with a single block.	Intraarticular injection of local anesthetic and steroid in 29 patients	Radiofrequency neurotomy for 90 seconds at 80°C in 27 patients	Roland-Morris questionnaire, Visual Analog Scale, Oswestry Disability Index, analgesic intake	6 months	Pain relief and functional improvement were observed in both groups. There were no significant differences between the 2 groups for pain relief and functional status improvement.	Lack of placebo group. Relatively short-term follow-up.	Randomized, double-blind trial with single diagnostic block with intraarticular injection	Both groups showed improvement. Effectiveness of both modalities at 6 months in both groups.

Table 9 (cont.). Study characteristics of randomized controlled trials and observational studies assessing lumbar radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
Yun et al. 2012 (124) Randomized, active controlled trial Quality Scores: Cochrane = 9/12 IPM-QRB = 26/48	57 patients with facet syndrome were assigned to 2 groups with 32 patients in the fluoroscopy group and 25 patients in the under ultrasonography group without diagnostic blocks	Intraarticular injection of lidocaine and triamcinolone under fluoroscopic guidance	Intraarticular injection of lidocaine and triamcinolone under ultrasonic guidance	Visual Analog Scale, physician's and patients' global assessment (PhyGA, PaGA), modified Oswestry Disability Index	One week, one and 3 months	Each group showed significant improvement from the facet joint injections. However at one week, one month, and 3 months after injections, no significant differences were observed between the groups.	Randomized trial	Short-term follow-up with no diagnostic blocks, thus increasing the potential for inclusion of patients without facet joint pain. The aim of study mainly was to confirm if ultrasonic imaging was appropriate.	The study showed positive results in both groups with intraarticular steroid injections with a short-term follow-up whether performed under ultrasonic guidance or fluoroscopy.

there were no trials available for thoracic intraarticular injection therapy.

The evidence for lumbar facet joint neurotomy is Level I for short-term effectiveness (< 6 months) and Level II for long-term effectiveness of 6 months or longer based on 8 moderate to high quality trials showing effectiveness (108,113,115,116,117,118,122,123), and one moderate to high quality trial (114) showed a lack of effectiveness. These results are in agreement with previously published systematic reviews (55-61). With recent publications, significant evidence has been demonstrated for lumbar radiofrequency neurotomy; however, multiple studies continue to include a small number of patients often with short-term follow-up. Among the long-term trials with effectiveness assessed at least at one year, Civelek et al (117) included 50 patients. Tekin et al (116) included 20 patients in the conventional radiofrequency neurotomy group; whereas, van Kleef et al (115) included only 15 patients in the radiofrequency neurotomy group showing positive results with a total number of 85 patients included. In contrast, van Wijk et al (114) showed a lack of effectiveness in 40 patients undergoing radiofrequency neurotomy.

The other studies by Cohen et al (108) included 14 patients with dual blocks; Nath et al (113) included 20 patients; Dobrogowski (118) included 45 patients; Moon et al (122) included 82 patients utilizing 2 different types of techniques; and Lakemeier et al (123) included 27 patients. Even though meta-analysis was not feasible based on a lack of homogeneity, others have attempted meta-analysis (59). Overall, other studies also showed similar results. Poetscher et al (59) and Leggett et al (58) showed positive results including the same studies included in this systematic review with best evidence synthesis. Both systematic reviews concluded that radiofrequency neurotomy has significant efficacy; however, others have showed a lack of efficacy (66,67). Overall, multiple deficiencies in these systematic reviews have been pointed out, including the small sample size of patients and the lack of homogeneity. Saltychev and Laimi (156) criticized the systematic review by Poetscher et al (59) for its lack of homogeneity. They noted that all conclusions were drawn from 2 studies (114,116), which were responsible for 85% of the entire synthesis. Further, the trial by Leclaire et al (165) was utilized in all recent systematic reviews (58,59,66,67), which has been considered as inappropriate and was excluded from this systematic review since the authors themselves have acknowledged multiple



Table 10. Study characteristics of randomized trials and observation studies assessing cervical and thoracic radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>CERVICAL RADIOFREQUENCY</b>									
Lord et al., 1996 (128) Randomized, sham control, double-blind Quality Scores: Cochrane = 11/12 IPM-QRB = 45/48	24 patients selected in a specialty cervical spine research unit in Australia suffering with chronic pain of cervical facet joint origin after whiplash injury and have failed conservative management. The diagnosis was confirmed with the use of double-blind, placebo-controlled local anesthetics with complete pain relief.	Sham control with placement of the needles with injection of local anesthetic without radiofrequency neurotomy.	Radiofrequency group 90 second lesion at 80°C of medial branch; control group received sham treatment with electrode insertion.  Even though this is described as a placebo-control treatment, maybe it is better to be called sham control since each patient received local anesthetic block which has been shown to be effective in itself (83).  Authors also produced multiple lesions at each level.	0 to 5 of 100 on visual analog scale; word count 3 or less on McGill Pain questionnaire.	3, 6, and 12 month follow-up	Median time to return of pain in treatment group was 263 days; 8 days in control group; 10 patients underwent second procedures with varying results.	Highly controlled design with meticulous diagnostic techniques and radiofrequency neurotomy.	Small group size and has been criticized for creative statistical analysis.	Efficacy was shown even though the study has been criticized for small group size and variations with creative statistical analysis. This study continues to be the landmark evaluation to show efficacy of radiofrequency neurotomy in the cervical spine.
Sapir & Gorup, 2001 (159) Prospective Quality Score: IPM-QRBNR = 40/48	32 litigants and 18 non-litigants underwent radiofrequency neurotomy.  Patients with cervical whiplash who remained symptomatic after 20 weeks of conservative management were included.  Inclusion criteria were 80% reduction in pain with controlled comparative local anesthetic blocks.  50 patients underwent radiofrequency neurotomy and 46 patients completed the study.	No control available.	The details for radiofrequency neurotomy were not provided.	Visual analog scale and self-report of improvement.		66% of the patients in the litigation group and 71% of the patients in the non-litigation group reported relief for more than one year. Time to recurrence defined as 50% return of pain was 8.0 ± 2.0 months. The frequency of recurrence of pain was similar in both groups.	Appropriate selection criteria with outcomes assessment in a prospective study.	Nonrandomized study with rather small number of patients.	The results were positive in this study in both litigants and non-litigants; however, there were only 32 litigants and 18 non-litigants undergoing radiofrequency neurotomy. Further, the difference between groups in the degree of symptomatology or response to treatment did not reach significance.

Table 10 (cont.). Study characteristics of randomized trials and observation studies assessing cervical and thoracic radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<p>MacVicar et al, 2012 (160)</p> <p>Prospective</p> <p>Quality Score: IPM-QRBNR = 38/48</p>	<p>A total of 104 patients selected on the basis of complete relief of pain following controlled, diagnostic, medial branch blocks were treated with radiofrequency neurotomy. The study was performed at 2 centers in New Zealand. A heterogeneous group of patients were included with suspicion of cervical zygapophysial joint pain. Patients were selected following the controlled comparative local anesthetic blocks with 100% pain relief concordant with duration of local anesthetic.</p>	<p>No control available.</p>	<p>Radiofrequency neurotomy was performed by placing the needles parallel to the medial branches, with creation of sufficient lesions in the sagittal plane, with 16-gauge 10 cm electrodes with 5 mm exposed tips. Radiofrequency was performed at 80°C or 85°C for 90 seconds for each lesion.</p>	<p>Successful outcome was defined as complete relief of pain, or at least 80% relief, for at least 6 months, with complete restoration of activities of daily living, no need for any further health care, and return to work.</p>		<p>In the 2 practices, 74% and 61% of the patients achieved a successful outcome. Relief lasted 17 to 20 months from the first radiofrequency neurotomy and 15 months for repeat treatments. Patients maintained relief for a median duration of 20 to 26 months, with 60% still having relief at follow-up.</p>	<p>The rigorous study was performed utilizing a rigorous criteria in a practical setting in New Zealand with impressive results.</p>	<p>Observational study performed in 2 different practices.</p>	<p>Positive results in a long-term follow-up with strict inclusion criteria with a meticulous technique with impressive results.</p>
<p>Speldewinde, 2011 (161)</p> <p>Prospective</p> <p>Quality Score: IPM-QRBNR = 39/48</p>	<p>A total 151 procedures were performed in the cervical spine on 130 patients. During the period from 2001 to 2010, patients were selected for radiofrequency thermal neurotomy in whom a diagnosis of cervical zygapophysial joint pain had been established with at least 2 fluoroscopically guided diagnostic medial branch nerve or intraarticular injections providing at least 80% relief in the index pain for the duration of action of local anesthetic used.</p>	<p>No control available.</p>	<p>Radiofrequency was performed at 80°C for 90 seconds for medial branches.</p>	<p>Numeric Rating Scale, Functional Rating Index, Activities of Daily Living, General Health Questionnaire, psychiatric morbidity</p>		<p>Cervical radiofrequency neurotomy was successful in 76% of the patients. The outcomes were similar in all 3 regions. A significant proportion of patients had relief for longer than one year. Average pain relief was 12 months in the cervical spine with average of 88% pain relief.</p>	<p>Even though study was prospective, design was appropriate and strict inclusion criteria with meticulous technique were utilized. Excellent outcome measures</p>	<p>Observational study.</p>	<p>Positive results. The study was performed in a community setting giving more of a practical setting in Australia.</p>

Table 10 (cont.). Study characteristics of randomized trials and observation studies assessing cervical and thoracic radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Study Characteristic Methodological Quality Scoring	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>CERVICAL FACET JOINT NERVE BLOCKS</b>										
Manchikanti et al, 2010 (83)	Randomized, double-blind, active-control	120 patients were recruited from consecutive new patients presenting to an interventional pain management practice with neck pain without suspected disc herniation or radiculitis. All the patients had conservative management and were judged to be positive for facet joint pain utilizing controlled comparative local anesthetic blocks with 80% pain relief and ability to perform previously painful movements. 60 patients received local anesthetic with steroid and another 60 patients received local anesthetics alone. 30 patients in each group also received Sarapin with local anesthetic and steroids.	Active control with local anesthetic and Sarapin	Cervical medial branch nerve blocks with fluoroscopy were performed utilizing local anesthetic with or without Sarapin or steroid.	Measured numeric pain scores, Neck Pain Index, opioid intake, and employment status at baseline. The procedures were repeated upon the return of pain and deterioration in functional status to less than 50%.	3, 6, 12, 18, and 24 months	85% of the patients with local anesthetic only and 92% of the patients with steroid reported significant pain relief at 12 months, whereas the statistics were 85% in local anesthetic group and 93% in steroid group at the end of 2 years. Functional status improvement of 50% or more by Neck Disability Index was seen in 63% and 68% at 12 months and 70% and 75% at 24 months in local anesthetic group and steroid group.	Randomized trial with relatively large proportion of patients with 2-year follow-up, with inclusion of patients diagnosed with controlled diagnostic blocks.	Lack of placebo group	This is the first study conducted evaluating therapeutic medial branch blocks in a randomized double-blind fashion, with effectiveness of facet joint nerve blocks with or without steroids.
Manchikanti et al, (152)	Prospective	100 consecutive patients meeting the diagnostic criteria of facet joint pain by means of comparative, controlled diagnostic blocks, with disabling chronic neck pain of various origins of at least 6 months duration and have failed conservative management were included.	No control available.	Medial branch blocks with fluoroscopy with bupivacaine with or without methylprednisolone. Patients had repeat blocks as clinically indicated.	Pain relief, Oswestry Disability Index, psychological status, work status	Timing: 3 months, 6 months, and 12 months	Significant pain relief at 3, 6, and 12 months, compared to baseline measurements. There was also significant improvement in disability status, psychological status, and return to work. Significant pain relief was observed at 92% at 3 months, 82% at 6 months, and 56% at 12 months.	First trial available	Non-randomized	Positive This was the first evaluation ever published in the cervical spine evaluating the role of therapeutic cervical medial branch blocks.
	Quality Score: IPM-QRBNR = 37/48									

Table 10 (cont.). Study characteristics of randomized trials and observation studies assessing cervical and thoracic radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Study Characteristic	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>CERVICAL INTRAARTICULAR INJECTIONS</b>										
Barnsley et al, 1994 (119)	Randomized, double-blind, active-control	41 patients with involvement of one or more cervical zygapophysial joints after automobile accidents with median duration of pain of 39 months were randomly assigned into 2 groups.	Active control trial with injection of intraarticular local anesthetic.	Intraarticular injection of 5.7 mg betamethasone or 1 mL intraarticular bupivacaine	Pain relief		No significant difference in duration of pain relief. Median duration of time to return of pain to 50% was 3 days in the steroid group and 3.5 days in the local anesthetic group.	Randomized controlled trial.	Small sample size, extremely short follow-up period.	Lack of effectiveness. Authors injected local anesthetic or steroid into the joint, thus this is not placebo controlled, it is rather an active-control trial.
Park & Kim, 2012 (129)	Randomized, active control	200 patients were studied in each group either with intraarticular injections or conservative management. Patients were selected for therapeutic intraarticular injections if they were positive for facet joint pain utilizing dual diagnostic blocks with concordant 80% pain relief.	Conservative management controlled trial.	Intraarticular injections were performed with 0.5 mL of 1% lidocaine and 5 mg of triamcinolone and 187.5 international units of hyaluronidase. Patients also received either onabotulinumtoxinA or trigger point injections if they required; however, these were of a small number.	Cervical range of motion, numeric rating scale for pain, comorbid tension type headache		Patients receiving intraarticular injections on one occasion showed increased cervical range of motion, increased mean numeric rating scale pain reduction, and decreased incidence of combined tension-type headache compared with control group receiving conservative management during follow-up.	Randomized controlled trial in a large proportion of patients.	Intraarticular injections were compared with conservative management. The study was also confounded with multiple deficiencies including trigger point injections and Botox injections in some patients with greater than 20% withdrawal rate.	Undetermined This study showed effectiveness of intraarticular injections.

Table 10 (cont.). Study characteristics of randomized trials and observation studies assessing cervical and thoracic radiofrequency neurotomy, facet joint nerve blocks, and intra-articular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>THORACIC RADIOFREQUENCY</b>									
Joo et al, 2013 (120) Randomized, double-blind, active control Quality Scores: Cochrane = 10/12 IPM-QRB = 38/48	40 patients with recurrent thoracolumbar facet joint pain after successful thermal radiofrequency ablation (RFA) defined as a numeric rating scale (NRS) score of 7 or a revised Oswestry disability index (ODI) of 22% were randomly allocated to 2 groups receiving either the same repeated RFA (n = 20) or alcohol ablation (AA) (n = 20).	Active control with alcohol.	Patients were provided with similar interventions with placement of radiofrequency needles, electric stimulation, contrast medium injection, local anesthetic injection followed by either radiofrequency thermoneurolysis for 90 seconds at 90°C or injection of 1 mL volume over a period of 15 seconds.	The recurrence rate was assessed with NRS and ODI and adverse events		After RFA and AA, one and 17 patients, respectively, were without recurring thoracolumbar facet joint pain. The median effective periods in the RFA and AA groups were 10.7 (range 5.4–24) and 24 (range 16.8–24) months, respectively (P = 0.000).	Randomized, double-blind, active control trial with appropriate outcomes assessment.	Small sample size. Selection criteria which included only the patients who required repeat thoracolumbar facet joint neurotomy after prior successful procedure.	This trial is the first of its nature for the thoracic spine in a randomized fashion with active control design. Specific importance is that they selected only the patients who had responded successfully with the first radiofrequency treatment for at least 6 months and then randomized them to assess the differences between alcohol injection and radiofrequency neurotomy.  Alcohol treatment was superior to radiofrequency for recurrent pain; however, this also shows effectiveness of radiofrequency neurotomy though inferior to alcohol with long-term follow-up of 24 months.

Table 10 (cont.). Study characteristics of randomized trials and observation studies assessing cervical and thoracic radiofrequency neurotomy, facet joint nerve blocks, and intraarticular injections.

Study	Number of Patients & Selection Criteria	Control	Interventions	Outcome Measures	Time of Measurement	Results	Strengths	Weaknesses	Conclusions
<b>THORACIC/FACET/JOINT NERVE BLOCKS</b>									
Manchikanti et al, 2012 (84) Randomized, double-blind, active controlled trial Quality Scores: Cochrane = 10/12 IPM-QRB = 45/48	100 patients were included with 50 patients in each of the local anesthetic and steroid groups. Selection was with controlled diagnostic blocks with criterion standard of 80% concordant pain relief.	Local anesthetic only.	Local anesthetic patients received thoracic medial branch blocks with bupivacaine. Local anesthetic with steroid patients received thoracic medial branch blocks with bupivacaine and non-particulate betamethasone.	Numeric pain scores, Oswestry Disability Index, opioid intake, and return to work status. Significant pain relief was defined as > 50% relief. Significant functional improvement was > 40% reduction of Oswestry Disability Index.	All outcomes were assessed at baseline, 6 months, 12 months, and 24 months.	In local anesthetic group, 80% of patients showed significant pain relief and functional improvement at 12 and 24 months. In local anesthetic with steroid group, 84% of patients showed significant pain relief and functional improvement at 12 months and 24 months. The majority of patients experienced significant pain relief for 46 to 47 weeks, requiring approximately 3 to 4 treatments with an average relief of 14 to 16 weeks per episode of a treatment.	Significant pain relief was shown in 85% in local anesthetic group and 90% in local anesthetic with steroids group at the end of the 2 year study period with controlled diagnostic blocks	Randomized trial with relatively large proportion of patients with 2-year follow-up, with inclusion of patients diagnosed with thoracic blocks	The majority of patients in both groups experienced significant pain relief and improvement in functional status. Therapeutic thoracic medial branch blocks, with or without steroid.
Manchikanti et al, 2006 (162) Prospective outcome study Quality Score: IPM-QRBNR = 37/48	55 consecutive patients, all meeting diagnostic criteria for thoracic facet joint pain	None	Thoracic facet joint nerve blocks performed using bupivacaine with or without Sarapin and depomethylprednisolone	Measured numeric pain scores, Oswestry Disability Index, employment status, and Pain Patient Profile.	3, 6, 12, 24, and 36 months.	Significant (≥ 50%), was observed in 71% of the patients at 3 months and 6 months, 76% at 12 months, 71% at 24 months, and 69% at 36 months.			Therapeutic thoracic medial branch blocks were an effective modality of treatment in managing chronic thoracic pain secondary to facet joint involvement confirmed by controlled, comparative local anesthetic blocks. Positive short-term and long-term relief.

Effectiveness of Therapeutic Facet Joint Interventions

Table 11. *Efficacy of lumbar radiofrequency, facet joint nerve blocks, and intraarticular injections.*

Study Study Characteristic Methodological Quality Scoring	Patients	Interventions	Pain Relief and Function			Results			Comments
			3 mos.	6 mos.	12 mos.	Short-Term ≤ 6 mos.	Long-Term		
							> 6 mos.	≥ 1 year	
<b>LUMBAR RADIOFREQUENCY NEUROTOMY</b>									
Civelek et al, 2012 (117)  RA, AC  Quality Scores: Cochrane = 9/12 IPM-QRB = 28/48	100	CRF = 50 Facet joint nerve blocks = 50	NA	92% vs. 75%	90% vs. 69%	NA	P	P	Effective for short and long-term
Cohen et al, 2010 (108)  RA, DB  Quality Scores: Cochrane = 8/12 IPM-QRB = 28/48	“0” block = 51 One block = 20 Two blocks = 14	CRF	“0” group = 33% One block = 39% Two blocks = 64%	NA	NA	P in dual block group	NA	NA	Effective in short-term results with application of dual blocks  Not effective with no or single diagnostic blocks.
Nath et al, 2008 (113)  RA, DB, Sham control  Quality Scores: Cochrane = 12/12 IPM-QRB = 42/48	40	Radiofrequency = 20  Sham = 20	NA	Significant proportion of patients in interventional group	NA	P for radiofrequency  N for sham or active	P for radiofrequency  N for sham or active	NA	Effective for short and long-term
Tekin et al, 2007 (116)  RA, AC and sham, DB  Quality Scores: Cochrane = 12/12 IPM-QRB = 37/48	60	CRF = 20 PRF = 20 Control = 20	NA	SI with CRF	SI with CRF	NA	P for radiofrequency  N for sham	P for radiofrequency  N for sham	Effective in long-term
van Wijk et al, 2005 (114)  RA, DB, Sham control  Quality Scores: Cochrane = 12/12 IPM-QRB = 36/48	81	Radiofrequency = 40 Sham = 41	27.5% vs. 29.3%	27.5% vs. 29.3%	27.5% vs. 29.3%	N	N	N	Lack of effectiveness with short- and long-term

Table 11 (cont.). *Efficacy of lumbar radiofrequency, facet joint nerve blocks, and intraarticular injections.*

Study Study Characteristic	Patients	Interventions	Pain Relief and Function			Results			Comments
			3 mos.	6 mos.	12 mos.	Short-Term ≤ 6 mos.	Long-Term		
Methodological Quality Scoring						> 6 mos.	≥ 1 year		
Dobrogowski et al, 2005 (118)  RA, AC  Quality Scores: Cochrane = 10/12 IPM-QRB = 29/48	45	CRF	NA	60%	NA	NA	P	NA	Short- and long-term effectiveness
van Kleef et al, 1999 (115)  RA, DB, sham control  Quality Scores: Cochrane = 12/12 IPM-QRB = 40/48	31	Radiofrequency = 15 Sham = 16	60% vs. 25%	47% vs. 19%	47% vs. 13%	P for radiofrequency  N for sham or active	P for radiofrequency  N for sham	P for radiofrequency  N for sham	Effectiveness with short- and long-term
Moon et al, 2013 (122)  Prospective, RA, comparative study  Quality Scores: Cochrane = 9/12 IPM-QRB = 38/48	Total = 82  Tunnel vision approach group – 41 patients included and 34 patients analyzed.	Radiofrequency neurotomy distal approach	SI in both groups	SI in both groups	NA	P	P	NA	Short- and long-term effectiveness
Lakemeier et al (123)  RA, DB  Quality Scores: Cochrane = 9/12 IPM-QRB = 37/48	Total = 56  Steroid group = 29 patients Radiofrequency group = 27 patients	Intraarticular lumbar facet joint steroid injections compared to lumbar facet joint radiofrequency denervation	NA	SI in both groups	NA	P	P	NA	Short- and long-term effectiveness
<b>LUMBAR FACET JOINT NERVE BLOCKS</b>									
Civelek et al, 2012 (117)  RA, AC  Quality Scores: Cochrane = 9/12 IPM-QRB = 28/48	100	LA with steroid = 50 CRF = 50	NA	75% vs. 92%	69% vs. 90%	NA	P	P	Long-term effectiveness



Effectiveness of Therapeutic Facet Joint Interventions

Table 11 (cont.). *Efficacy of lumbar radiofrequency, facet joint nerve blocks, and intraarticular injections.*

Study Study Characteristic	Patients	Interventions	Pain Relief and Function			Results			Comments
			3 mos.	6 mos.	12 mos.	Short-Term ≤ 6 mos.	Long-Term > 6 mos.    ≥ 1 year		
Manchikanti et al, 2010 (82)  RA, DB, AC  Quality Scores: Cochrane = 11/12 IPM-QRB = 45/48	120	LA with steroid = 60 LA = 60	82% vs. 83%	93% vs. 83%	85% vs. 84%	P	P	P	Short- and long-term effectiveness
Manchikanti et al, 2001 (112)  RA, AC  Quality Scores: Cochrane = 6/12 IPM-QRB = 34/48	73	LA with steroid = 41 LA = 32	100% vs 100%	75% vs 80%	75% vs 80%	P	P	P	Positive short and long-term results
<b>LUMBAR INTRAARTICULAR INJECTIONS</b>									
Carette et al, 1991 (110)  RA, DB, PC or AC  Quality Scores: Cochrane = 11/12 IPM-QRB = 40/48	97	Methylprednisolone acetate =49  Isotonic saline =48 patients	33% vs. 42%	22% vs. 10%	NA	N	N	NA	Lack of effectiveness
Fuchs et al, 2005 (111)  R, DB, AC  Quality Scores: Cochrane = 8/12 IPM-QRB = 26/48	60	Hyaluronic acid versus glucocorticoid with 6 injections.	Significant proportion of patients	Significant proportion of patients	NA	U	U	NA	Effectiveness undetermined
Ribeiro et al, 2013 (121)  RA, DB, AC  Quality Scores: Cochrane = 10/12 IPM-QRB = 32/48	60	Intraarticular injection group = 31 Intramuscular steroid injection group = 29	52% vs 45%	55% vs 38%	NA	P	P	NA	Short- and long-term effectiveness

Table 11 (cont.). *Efficacy of lumbar radiofrequency, facet joint nerve blocks, and intraarticular injections.*

Study Study Characteristic	Patients	Interventions	Pain Relief and Function			Results			Comments
			3 mos.	6 mos.	12 mos.	Short-Term ≤ 6 mos.	Long-Term > 6 mos.      ≥ 1 year		
Yun et al, 2012 (124)  RA  Quality Scores: Cochrane = 9/12 IPM-QRB = 26/48	Total = 57  Fluoroscopy group = 32 Ultrasonography group = 25	Intraarticular injection of local anesthetic and steroid	SI in both groups	NA	NA	<i>P</i>	NA	NA	Short-term effectiveness
Lakemeier et al, 2013 (123)  RA, DB  Quality Scores: Cochrane = 9/12 IPM-QRB = 37/48	Total = 56  Steroid group = 29 patients Radiofrequency group = 27 patients	Intraarticular lumbar facet joint steroid injections compared to lumbar facet joint radiofrequency denervation	NA	SI in both groups	NA	<i>P</i>	<i>P</i>	NA	Short-and long-term effectiveness

RA = randomized; DB = double-blind; AC = active control; ST = steroid; LA = local anesthetic; SAL = saline; SI = significant improvement; P = positive; N = negative; NA = not applicable

flaws in performing their trial. Further, multiple issues related to radiofrequency neurotomy and exclusion of studies is based on an author's own criteria such, as by LeClaire et al (165).

The evidence for cervical radiofrequency neurotomy was derived from only one high quality randomized controlled trial with an extremely small sample size of patients (128). There were no other trials; consequently, it appears that the level of evidence of II or III may be appropriate. Since there was only one randomized double-blind controlled trial, it may be worthwhile to look at the evidence from nonrandomized prospective trials. The randomized trial by Lord et al (128) included 24 patients and compared percutaneous radiofrequency neurotomy to a sham procedure. Thus, the technique of radiofrequency thermoneurolysis was the same, but radiofrequency lesioning was not performed in the sham control group.

The inclusion criteria were rather strict with comparative local anesthetic blocks with 100% concordant pain relief. The post-treatment assessment was carried out with the Visual Analog Scale and the McGill Pain Questionnaire. The results showed one patient in the sham control group and 7 patients in the active treatment group were pain-free at the 27-week follow-up.

Their results also showed that the median time for return of pain to at least 50% of the preoperative level was 263 days in the active treatment groups; whereas it was 8 days in the sham control group. The authors concluded that this trial proved that radiofrequency neurotomy is capable of giving pain relief for up to and over a year. Even though this study was meticulously performed in an academic setting, it only included a small number of patients with whiplash injury and the technique with multiple lesions is not commonly utilized in the United States. This trial also faced significant criticism by Carragee et al (157) for multiple aspects that have not been widely recognized and criticized in the past, which included the differences in baseline characteristics of patients among both groups and the nature of blinding. In fact, Carragee et al (157) reported that the integrity of the blinding was in doubt related to the fact that 42% of the active group developed long-term anesthetic or dysesthetic areas of the skin, whereas none of the patients in the sham control group developed changes. Thus, Carragee et al (157) felt that there was no significant blinding technique applied with ability of the participants to recognize their group assignment. In addition, litigation also had no significant difference in the outcomes. Overall, the

Effectiveness of Therapeutic Facet Joint Interventions

Table 12. Results of randomized trials of cervical and thoracic radiofrequency neurotomy, facet joint, nerve blocks and intraarticular injections.

Study Study Characteristic Methodological Quality Scoring	Patients	Interventions	Pain Relief and Function			Results			Comments
			3 mos.	6 mos.	12 mos.	Short-Term ≤ 6 mos.	Long-Term > 6 mos.    ≥ 1 year		
<b>CERVICAL RADIOFREQUENCY</b>									
Lord et al, 1996 (128) RA, sham control, DB Quality Scores: Cochrane = 11/12 IPM-QRB = 45/48	24	Conventional RFTN 80°C, 90 seconds  Sham = 12 Intervention = 12	NA	One of sham 7 of active	58% in active treatment group	P	P	P	Short- and long-term effectiveness
Sapir & Gorup, 2001 (159)  Prospective Quality Score: IPM-QRBNR = 40/48	50	Conventional RFTN 80°C, 90 seconds  Litigants = 32 Non-litigants = 18	NA	NA	66% litigant  71% non-litigant	NA	NA	P	Long-term effectiveness
MacVicar et al, 2012 (160)  Prospective Quality Score: IPM-QRBNR = 38/48	104	Conventional RFTN 80°C, 90 seconds  2 practices	NA	NA	74% vs 61%	NA	NA	P	Long-term effectiveness
Speldewinde, 2011 (161)  Prospective Quality Score: IPM-QRBNR = 39/48	130	Conventional RFTN 80°C, 90 seconds	NA	NA	76%	NA	NA	P	Long-term effectiveness
<b>CERVICAL FACET JOINT NERVE BLOCKS</b>									
Manchikanti et al, 2010 (83) RA, DB, AC Quality Scores: Cochrane = 11/12 IPM-QRB = 45/48	120	Local anesthetic = 60 Local anesthetic with steroid = 60	83% versus 85%	87% versus 95%	85% versus 92%	P	P	P	Short- and long-term effectiveness
Manchikanti et al, 2004 (152)  Prospective Quality Score: IPM-QRBNR = 37/48	100	Therapeutic medical branch blocks	92%	82%	56%	P	P	P	Long-term effectiveness
<b>CERVICAL INTRAARTICULAR INJECTIONS</b>									
Barnsley et al, 1994 (119) RA, DB, AC Quality Scores: Cochrane = 12/12 IPM-QRB = 36/48	41	LA = 20 Steroid = 21	20%	20%	20%	N	N	N	Lack of effectiveness

Table 12 (cont.). Results of randomized trials of cervical and thoracic radiofrequency neurotomy, facet joint, nerve blocks and intraarticular injections.

Study Study Characteristic Methodological Quality Scoring	Patients	Interventions	Pain Relief and Function			Results			Comments
			3 mos.	6 mos.	12 mos.	Short-Term ≤ 6 mos.	Long-Term		
							> 6 mos.	≥ 1 year	
Park & Kim, 2012 (129) RA, AC  Quality Scores: Cochrane = 6/12 IPM-QRB = 35/48	306	Non-injection group = 151 Nerve blocks = 155	U	U	U	U	U	U	Unable to determine effectiveness
<b>THORACIC RADIOFREQUENCY</b>									
Joo et al, 2013 (120)  Quality Scores: Cochrane = 10/12 IPM-QRB = 38/48	40	Radiofrequency neurotomy = 20 Alcohol injection = 20	SI in both groups	SI in both groups	SI in both groups	P	P	P	Short- and long-term effectiveness of radiofrequency neurotomy in alcohol injection group.
<b>THORACIC FACET JOINT NERVE BLOCKS</b>									
Manchikanti et al, 2012 (84) RA, DB  Quality Scores: Cochrane = 10/12 IPM-QRB = 45/48	100	Local anesthetic = 50 Local anesthetic with steroid = 50	79% vs 83%	79% vs 81%	80% vs 83%	P	P	P	Short- and long-term effectiveness

RA = randomized; DB = double-blind; P = prospective; R = retrospective; vs = versus; P = positive

results showed that in the sham control group 58% of the patients and in the active treatment group 25% of the patients had their pain come back immediately after the procedure at the 3-month follow-up. Lord et al (128) were obviously unable to avoid this issue of different return of pain in 2 different groups and this is a problem with any of the interventional techniques. In contrast to the criticism of Carragee et al (157), Dreyfuss and Baker (158) supported Lord et al's manuscript (128) for maintaining appropriate blinding of patients based on the fact that it was very difficult to maintain a lack of anesthetic effect and also the numerous difficulties encountered in performing such studies, which is evidenced by the lack of such studies thus far in the cervical spine.

The remaining radiofrequency neurotomy studies were observational in nature. Sapir and Gorup (159), in a 2001 study evaluated the effectiveness of radiofrequency medial branch neurotomy of cervical facet joints after whiplash injury with chronic neck pain in a design which compared litigants to nonlitigants. The inclusion criteria included involvement in a motor vehicle

accident at least 20 weeks prior to the study, failure to respond to conservative treatment, and a positive response to controlled, comparative, local anesthetic blocks. They included 50 patients who met inclusion criteria with at least 80% pain relief from comparative local anesthetic blocks and subsequently underwent radiofrequency neurotomy. However, only 46 of the patients completed the study with 29 in the litigation group (63%) and 17 in the nonlitigation group (37%). Subsequent to radiofrequency neurotomy, 21 patients, 14 patients in the litigation group and 7 in the nonlitigation group, experienced recurrence of pain within one year, whereas, 25 patients, 15 in the litigation group and 20 in the nonlitigation group, remained asymptomatic at the end of the one-year follow-up period. They showed that the return of pain, which they defined as 50% of pain returning, was approximately  $8.3 \pm 2.3$  months in the 21 patients whose pain returned within one year. There were no significant differences in relation to the outcomes between the litigant and nonlitigant groups. Overall the authors concluded that cervical radiofrequency neurotomy of facet joints in

chronic neck pain secondary to whiplash injury was an effective modality independent of litigation.

In another study by MacVicar (160), which was derived from 2 practices in New Zealand, a successful outcome was reported in 74% and 61% of patients with long-lasting relief of 17 to 20 months from the first radiofrequency neurotomy, and 15 months for repeat radiofrequency neurotomy. Considering the need for repeat treatments, which were provided appropriately, overall the patients maintained relief for a median duration of 20 to 26 months, with 60% continuing to have relief with one radiofrequency neurotomy procedure. The authors concluded that radiofrequency neurotomy is an effective technique when performed in a rigorous manner with appropriate selection of patients and consideration of the procedural requirements in chronic neck pain secondary to cervical facet joint involvement temporarily, but completely, relieved of pain, restoring patients to desired activities of daily living. In another study by Speldewinde (161), with 379 procedures, 272 or 72% of the procedures were considered successful by the patients, irrespective of the region treated. He showed a large effect size with significant improvement. He also showed that repetition of the procedure was highly successful. He concluded that radiofrequency neurotomy of not only the cervical facet joints, but also thoracic and lumbar facets and sacroiliac joints were uniformly successful with 72% of recipients obtaining an average of 86% reduction in pain for a period of 12 months. Other studies by McDonald et al (163) and Barnsley et al (164) also demonstrated significant progress on a long-term basis. McDonald et al (163), in an assessment of long-term follow-up of patients, performed cervical radiofrequency neurotomy for chronic neck pain and showed successful results with complete pain relief in 71% of the patients after an initial procedure; however, the pain returned after 290 days when failures were included. Otherwise, they reported 422 days of relief with all successful cases. The major deficiency was that it included only 28 patients, which appears to have been replicating the results of the randomized controlled trial (128) with very similar outcomes. Barnsley et al (164) also assessed percutaneous radiofrequency neurotomy for chronic neck pain in 35 patients with 47 procedures. The results showed patients receiving 36 procedures achieved 80% significant pain relief with a mean duration of pain relief of 36 weeks, with repeat procedures usually achieving reproducible pain relief.

For thoracic radiofrequency neurotomy the level of

evidence is Level III based on one high quality randomized controlled trial (120).

The level of evidence for facet joint nerve blocks is Level II in the lumbar, cervical, and thoracic regions based on 4 high quality randomized controlled trials (82-84,117) and one moderate to high quality randomized controlled trial (112). Further, there were no trials showing a lack of effectiveness.

Therapeutic facet joint nerve blocks were studied in all 3 regions with 2 high quality randomized controlled trials (82,117) in the lumbosacral region, and one moderate to high quality randomized controlled trial (112). One high quality randomized controlled trial in the thoracic (84) and cervical region (83). Four of the 5 manuscripts were from the same group of authors (82-84,112). In 2 manuscripts (82,83), 120 patients were included in each of the studies of lumbar and cervical facet joint pain (82,83); in studying thoracic facet joint pain (84) 100 patients were included; whereas, in one lumbar trial (112) 73 patients were included. The patients were all drawn from an interventional pain management practice. They had all failed conservative management and were judged to be positive for facet joint pain utilizing controlled comparative local anesthetic blocks with 80% pain relief as the criterion standard with ability to perform previously painful movements. In each group, an equal number of patients were allocated to receive either local anesthetic alone or local anesthetic with steroid. Outcome parameters included pain relief criteria and disability criteria with follow-ups at 3, 6, 12, 18, and 24 months. Significant pain relief was defined as greater than 50% relief with significant improvement in functional status of greater than 40%. In the lumbar region (82), the results showed significant pain relief in 85% of the patients receiving local anesthetic and 90% of the patients receiving local anesthetic with steroids at the end of the 2 year study period with an average of 5 to 6 total treatments. In the cervical spine (83), 85% of the patients in the local anesthetic only group and 93% of the patients in the steroid and local anesthetic group had significant improvement with a total of 5 to 6 procedures over a period of 2 years.

In the thoracic spine, the results were similar to the cervical and lumbar spines with 80% of the patients in the local anesthetic group and 84% of the patients in the local anesthetic and steroid group showing significant improvement at the end of 2 years with a total of 5 to 6 procedures. The fourth study, by Civelek et al (117), studied 100 patients with chronic low back pain who

failed conservative therapy and implemented strict selection criteria even though no diagnostic blocks were utilized. They used lumbosacral facet joint nerve blocks as the control group, whereas, the second group received conventional radiofrequency neurotomy. They followed the patients for 6 months and 12 months. At the end of one-year, 69% of the patients in the facet joint nerve block group showed significant improvement compared to 90% in the radiofrequency neurotomy group. Overall it showed the effectiveness of lumbar facet joint nerve blocks even though they were inferior to radiofrequency neurotomy. The final moderate to high quality study by Manchikanti et al (112), included 73 patients and compared a combination of bupivacaine with Sarapin and bupivacaine with Sarapin with the addition of steroids and showed positive results on a long-term basis.

These results are similar to some previously published systematic reviews (55-57); however, there were no other systematic reviews which appropriately studied the role of facet joint nerve blocks.

The evidence for the lumbar intraarticular injection of steroids is Level III, based on 3 high quality randomized controlled trials, showed effectiveness with short-term follow-up of less than 6 months (121,123,124); however, the results were also opposed by 2 moderate to high quality randomized controlled trials showing a lack of effectiveness (110,111). This level of evidence is similar to other published systematic reviews. In the cervical spine there were 2 randomized controlled trials of intraarticular injections (119,129) yielding evidence of Level III with one trial showing a lack of effectiveness and the second one showing undetermined results. These results are also similar to previous systematic reviews. There were no studies on intraarticular injections in the thoracic spine.

The disadvantages of this systematic review include the lack of metaanalysis; however, there was no clinical homogeneity among the trials. Further, it would be inappropriate to perform a systematic review based on some hypothetical principle if the trials are not clinically homogenous. Consequently, a best evidence synthesis appears to be the most appropriate in this setting. Other disadvantages include the continued paucity of literature about facet joint nerve blocks in all 3 regions and radiofrequency neurotomy in the cervical and thoracic regions as well as intraarticular injections in the cervical and thoracic regions. Future trials must be of appropriate size, draw from a population from practical settings, with a minimum long-term follow-up of one-year. Mul-

iple other issues related to facet joint interventions include placebo response, nocebo response, the role of sham procedures, technical aspects in performing a procedure, and finally the role of local anesthetic alone compared to steroids with sodium chloride solution or steroids with local anesthetic (165-224). Based on the present evidence, there is no additional effectiveness beyond the relief provided by local anesthetic blocks with the addition of steroids, bupivacaine specifically, in facet joint nerve blocks (82-84,112,152,162,180).

The rationale for intraarticular injections comes from steroids being used for treating inflammation. The literature abounds with reports that epidural corticosteroid injections have significant efficacy for their anti-inflammatory, immuno-suppressive, anti-edema effects and inhibition of neurotransmission within the C fibers (185-198). The same is supported with facet joint nerve blocks; with long-term symptomatic improvement very similar to the addition of steroids and even better than with steroids (82-84,112,152,162,180,189,190,225-234). The experimental evidence also shows a lack of effectiveness of adding steroids (189,190).

It has been postulated that local anesthetics provide relief by suppressing nociceptive discharge (190), blocking axonal transport (191,192), blocking the sympathetic reflex arc, blocking sensitization (193,194), and by their anti-inflammatory effects (195). Local anesthetics have been reported to have long-term effectiveness following local anesthetic nerve blocks or epidural injections (82-84,110,112,152,180,206-208,225-234).

The lack of placebo in active control trials is a major misunderstanding and a limitation. However, placebo control has been misunderstood in many cases. The reviewers have considered a local anesthetic injection as a placebo control. It is a well known fact that placebo control in any neural blockade is a difficult task. Further, it also adds ethical issues and difficulty with recruitment in the United States. However, multiple investigations performed in interventional pain management with descriptions of placebo control have design flaws (62,65,180,199-204). A solution's effect when injected into a closed space has been inappropriately appraised. Carrette et al (110,196) reported that the response is similar whether an injectate has a sodium chloride solution or a local anesthetic with a steroid. The response to both injections in both the intraarticular and epidural space was low. Thus, their study (110) shows that sodium chloride solution injected into an intraarticular space has similar effects as local anesthetic with a steroid; the conclusion is that intraarticular steroids are

not an effective therapy. The issue is also exemplified by Birkenmaier et al (205), utilizing either pericapsular injections or medial branch blocks, who then went on to perform cryoneurolysis. Not surprisingly, the results were superior in patients who were diagnosed using medial branch blocks rather than pericapsular injections of local anesthetic. This study was the basis for Chou and Huffman (7) to reject diagnostic lumbar facet joint nerve blocks as having value. Also, there are reports of different effects from different solutions such as local anesthetic, normal saline, and dextrose; the same is true when a solution is injected into the disc, facet joint, or multifidus muscle (206-213). It has been shown that a small volume of local anesthetic or normal saline abolishes muscle twitch induced by a low current (0.5 mA) during electrode location (206-209). Further, there is direct evidence for spinal cord involvement in placebo analgesia (210). It also has been shown that epidurally administered sodium chloride solution provides significant improvement in pain and function (196,214-217). Therefore, it can be concluded that local anesthetic's effect on cervical facet joint nerve blocks is not due to the placebo effect, even though some have mistakenly misinterpreted this to be the case (200,201,217,218).

Placebo effects are not expected to be seen in a high proportion of patients, nor are they expected to be long lasting with repeat interventions over a period of 2 years. However, the limitations of the lack of placebo must not be underestimated. If feasible, a placebo-controlled study with appropriate design that includes not injecting the placebo solution over the facet joint nerves, and subsequent results, would be highly valid and provide conclusive knowledge on the issue of placebo-controlled blocks. The issues related to placebo have been discussed extensively in recent years ultimately leading to the opinion that the placebo effect is an inconsistent measure in clinical studies, unless it is designed appropriately (166,167,169,170,219-224).

Another issue is related to the reliability of controlled, comparative local anesthetic blocks, which have been criticized, and their validity as precision diagnostic techniques has been questioned and debated (7,10,16-19,62,199,235-239). The issues related to the accuracy of diagnostic facet joint nerve blocks include the reference standard, prior exposure to opioids, sedation, systemic local anesthetic effect, and non-specific effect resulting in positive results (7,10,16-19,56,62,209,240-249). The validity of controlled facet joint nerve blocks as a gold standard or reference standard in the diagnosis of lumbar facet joint pain has been established (248,249).

A reference standard is established in surgical situations via biopsy or autopsy. However, these are difficult to apply in the diagnosis of chronic spinal pain of facet joint origin. Thus, the long-term or dedicated clinical follow-up of patients appears to be the only solution in establishing a reference standard with controlled facet joint nerve blocks (250). Based on the criterion standard of long-term follow-up, controlled diagnostic lumbar facet joint nerve blocks have been shown to be valid utilizing the criteria of 80% pain relief and the ability to perform previously painful movements, with a sustained diagnosis of lumbar facet joint pain in at least 89.5% of the patients at the end of 2-year follow-up (248). However, the diagnosis was sustained in only 51% of the patients with 50% relief at the end of 2 years (248). Thus, the controlled diagnostic blocks utilized in this study appear to be reliable.

#### **4.0 CONCLUSION**

This systematic review shows Level II evidence for long-term effectiveness of radiofrequency neurotomy in the lumbar and cervical spines, for facet joint nerve blocks in the cervical, thoracic, and lumbar spine, and Level III evidence for thoracic radiofrequency neurotomy, lumbar intraarticular injections and cervical intraarticular injections. This systematic review was performed utilizing strict inclusion criteria and methodological quality assessment criteria. Overall, the results appear to be somewhat superior in patients who receive conventional radiofrequency neurotomy after undergoing controlled diagnostic blocks.

#### **ACKNOWLEDGMENTS**

The authors wish to thank Vidyasagar Pampati, MSc, for statistical assistance, and Tom Prigge, MA, Laurie Swick, BS, and Sanjana Pampati, BS, for manuscript review, and Tonie M. Hatton and Diane E. Neihoff, transcriptionists, for their assistance in preparation of this manuscript. We would like to thank the editorial board of Pain Physician for review and criticism in improving the manuscript.

#### **Author Affiliations**

Dr. Manchikanti is Medical Director of the Pain Management Center of Paducah, Paducah, KY, and Clinical Professor, Anesthesiology and Perioperative Medicine, University of Louisville, Louisville, KY.

Dr. Kaye is Professor and Chair, Department of Anesthesia, LSU Health Science Center, New Orleans, LA.

Dr. Boswell is Professor and Chair, Department of

Anesthesiology and Perioperative Medicine, University of Louisville, Louisville, KY

Dr. Bakshi is President of Manhattan Spine and Pain Medicine, New York, NY.

Dr. Gharibo is Medical Director of Pain Medicine and Associate Professor of Anesthesiology and Orthopedics, Department of Anesthesiology, NYU Langone-Hospital for Joint Diseases, NYU School of Medicine, New York, NY.

Dr. Grami is an Interventional Pain Physician at Geisinger Medical Center Interventional Pain Center Woodbine, Danville, PA.

Dr. Grider is Medical Director, UK HealthCare Pain Services, Division Chief, Pain and Regional Anesthesia, and Associate Professor, Department of Anesthesiology, University of Kentucky, Lexington, KY.

Dr. Gupta is a Consultant in Pain Medicine and Anaesthesia, Bradford Teaching Hospital NHS Foundation Trust, Bradford Royal Infirmary, Bradford, United Kingdom.

Dr. Jha is an anesthesiology resident physician at Rush University Medical Center (RUMC), Chicago, IL.

Dr. Mann is Medical Director, Garden State Pain Management, Whiting, NJ.

Dr. Nampiaparampil is Assistant Professor, Rehabilitation Medicine, New York University School of Medicine, New York, NY.

Dr. Sharma is a Consultant in Pain Medicine, Department of Pain Medicine, The Walton Centre for Neurology and Neurosurgery NHS Foundation Trust, Liverpool, and a Fellow of the Faculty of Pain Medicine of the Royal College of Anaesthetists and Treasurer of British Pain Society Interventional Pain Medicine Special Interest Group, United Kingdom.

Dr. Shroyer is an Interventional Pain Physician at Florida Pain Management Associates, Sebastian, FL.

Dr. Singh is Medical Director, Spine Pain Diagnostics Associates, Niagara, WI

Dr. Soin is Medical Director, Ohio Pain Clinic, Centerville, OH.

Dr. Vallejo is Director of Research, Millennium Pain Center, Bloomington, IL; and Adjunct Professor of Biology, Illinois State University, Normal, IL

Dr. Wargo is an Interventional Pain Physician at the McFarland Clinic, Mary Greeley Medical Center, Ames, IA.

Dr. Hirsch is Vice Chief of Interventional Care, Chief of NeuroInterventional Spine, Service Line Chief of Interventional Radiology, Director of NeuroInterventional Services and Neuroendovascular Program, Massachusetts General Hospital; and Associate Professor, Harvard Medical School, Boston, MA.

**Conflict of interest:**

Dr. Manchikanti has provided limited consulting services to Semnur Pharmaceuticals, Incorporated, which is developing nonparticulate steroids.

Dr. Kaye is a speaker for Depomed, Inc.

Dr. Gupta has been paid honorarium for presenting at meetings and teaching on the interventional pain medicine cadaver courses and by pharmaceutical companies for presenting to health care professionals. Pharmaceutical companies and companies that manufacture equipments used in pain medicine have supported meetings organized by Dr S Gupta

Dr. Vallejo receives research support from Cephalon/Teva, BioDelivery Sciences International, Inc., Mundipharma Research GmbH & Co., AstraZeneca, Purdue Pharma, LP, and Theravance, and is a consultant for Nevro and Kimberly-Clark.

Dr. Hirsch is a consultant for Medtronic.



## REFERENCES

- US Burden of Disease Collaborators. The state of US health, 1999-2010: Burden of diseases, injuries, and risk factors. *JAMA* 2013; 310:591-608.
- Institute of Medicine (IOM). *Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research*. The National Academies Press, Washington, DC, June 29, 2011.
- Martin BI, Turner JA, Mirza SK, Lee MJ, Comstock BA, Deyo RA. Trends in health care expenditures, utilization, and health status among US adults with spine problems, 1997-2006. *Spine (Phila Pa 1976)* 2009; 34:2077-2084.
- Gaskin DJ, Richard P. The economic costs of pain in the United States. *J Pain* 2012; 13:715-724.
- Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, Williams G, Smith E, Vos T, Barendregt J, Murray C, Burstein R, Buchbinder R. The global burden of low back pain: Estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis* 2014; 73:968-974.
- Hoy D, March L, Woolf A, Blyth F, Brooks P, Smith E, Vos T, Barendregt J, Blore J, Murray C, Burstein R, Buchbinder R. The global burden of neck pain: Estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014; 73:1309-1315.
- Chou R, Huffman L. *Guideline for the Evaluation and Management of Low Back Pain: Evidence Review*. American Pain Society, Glenview, IL, 2009.
- Manchikanti L, Singh V, Falco FJE, Benyamin RM, Hirsch JA. Epidemiology of low back pain in adults. *Neuromodulation* 2014; 17:3-10.
- Manchikanti L, Pampati V, Falco FJE, Hirsch JA. Growth of spinal interventional pain management techniques: Analysis of utilization trends and Medicare expenditures 2000 to 2008. *Spine (Phila Pa 1976)* 2013; 38:157-168.
- Manchikanti L, Abdi S, Atluri S, Benyamin RM, Boswell MV, Buenaventura RM, Bryce DA, Burks PA, Caraway DL, Calodney AK, Cash KA, Christo PJ, Cohen SP, Colson J, Conn A, Corder HJ, Coubarous S, Datta S, Deer TR, Diwan SA, Falco FJE, Fellows B, Geffert SC, Grider JS, Gupta S, Hameed H, Hameed M, Hansen H, Helm II S, Janata JW, Justiz R, Kaye AD, Lee M, Manchikanti KN, McManus CD, Onyewu O, Parr AT, Patel VB, Racz GB, Sehgal N, Sharma M, Simopoulos TT, Singh V, Smith HS, Snook LT, Swicegood J, Vallejo R, Ward SP, Wargo BW, Zhu J, Hirsch JA. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain: Part II: Guidance and recommendations. *Pain Physician* 2013; 16:S49-S283.
- Rubinstein SM, van Middelkoop M, Assendelft WJ, de Boer MR, van Tulder MW. Spinal manipulative therapy for chronic low-back pain: An update of a Cochrane review. *Spine (Phila Pa 1976)* 2011; 36:E825-E846.
- van Middelkoop M, Rubinstein SM, Kuijpers T, Verhagen AP, Ostelo R, Koes BW, van Tulder MW. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J* 2011; 20:19-39.
- Manchikanti L, Pampati V, Falco FJE, Hirsch JA. An updated assessment of utilization of interventional pain management techniques in the Medicare population: 2000 - 2013. *Pain Physician* 2015; 18:E115-E127.
- Manchikanti L, Pampati V, Singh V, Falco FJE. Assessment of the escalating growth of facet joint interventions in the Medicare population in the United States from 2000 to 2011. *Pain Physician* 2013; 16:E365-E378.
- Manchikanti L, Helm II S, Singh V, Hirsch JA. Accountable interventional pain management: A collaboration among practitioners, patients, payers, and government. *Pain Physician* 2013; 16:E635-E670.
- Rubinstein SM, van Tulder M. A best-evidence review of diagnostic procedures for neck and low-back pain. *Best Pract Res Clin Rheumatol* 2008; 22:471-482.
- Hancock MJ, Maher CG, Latimer J, Spindler MF, McAuley JH, Laslett M, Bogduk N. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J* 2007; 16:1539-1550.
- Bogduk N. On diagnostic blocks for lumbar zygapophysial joint pain. *F1000 Med Rep* 2010; 2:57.
- Boswell MV, Manchikanti L, Kaye AD, Bakshi S, Gharibo CG, Gupta S, Jha S, Nampiaparampil DE, Simopoulos TT, Hirsch JA. A best-evidence systematic appraisal of the diagnostic accuracy and utility of facet (zygapophysial) pain injections in chronic spinal pain. *Pain Physician* 2015; \_\_\_\_\_.
- Kuslich SD, Ulstrom CL, Michael CJ. The tissue origin of low back pain and sciatica: A report of pain response to tissue stimulation during operation on the lumbar spine using local anesthesia. *Orthop Clin North Am* 1991; 22:181-187.
- Kalichman L, Li L, Kim DH, Guermazi A, Berkin V, O'Donnell CJ, Hoffmann U, Cole R, Hunter DJ. Facet joint osteoarthritis and low back pain in the community-based population. *Spine (Phila Pa 1976)* 2008; 33:2560-2565.
- Eubanks JD, Lee MJ, Cassinelli E, Ahn NU. Prevalence of lumbar facet arthrosis and its relationship to age, sex, and race: An anatomic study of cadaveric specimens. *Spine (Phila Pa 1976)* 2007; 32:2058-2062.
- Harris RI, Macnab I. Structural changes in the lumbar intervertebral discs; their relationship to low back pain and sciatica. *J Bone Joint Surg Br* 1954; 36-B:304-322.
- Bykowski JL, Wong WH. Role of facet joints in spine pain and image-guided treatment: A review. *AJNR Am J Neuroradiol* 2012; 33:1419-1426.
- Tessitore E, Molliqaj G, Schatlo B, Schaller K. Clinical evaluation and surgical decision making for patients with lumbar discogenic pain and facet syndrome. *Eur J Radiol* 2015; 84:765-770.
- Gellhorn AC, Katz JN, Suri P. Osteoarthritis of the spine: The facet joints. *Nat Rev Rheumatol* 2013; 9:216-224.
- Igarashi A, Kikuchi S, Konno S, Olmarker K. Inflammatory cytokines released from the facet joint tissue in degenerative lumbar spinal disorders. *Spine (Phila Pa 1976)* 2004; 29:2091-2095.
- Schulte TL, Filler TJ, Struwe P, Liem D, Bullman V. Intra-articular meniscoid folds in thoracic zygapophysial joints. *Spine (Phila Pa 1976)* 2010; 35:E191-E197.
- Kim JS, Kroin JS, Buvanendran A, Li X, van Wijnen AJ, Tuman KJ, Im HJ. Characterization of a new animal model for evaluation and treatment of back pain due to lumbar facet joint osteoarthritis. *Arthritis Rheum* 2011; 63:2966-2973.
- Masini M, Paiva WS, Araujo AS, Jr. Anatomical description of the facet joint innervation and its implication in the treatment of recurrent back pain. *J Neurosurg Sci* 2005; 49:143-146.
- Cavanaugh JM, Lu Y, Chen C, Kallakuri S. Pain generation in lumbar and cervical facet joints. *J Bone Joint Surg Am* 2006; 88:63-67.
- Bogduk N, Wilson AS, Tynan W. The human lumbar dorsal rami. *J Anat* 1982; 134:383-397.

33. Bogduk N. The clinical anatomy of the cervical dorsal rami. *Spine (Phila Pa 1976)* 1982; 7:319-330.
34. Ohtori S, Takahashi K, Chiba T, Yamagata M, Sameda H, Moriya H. Sensory innervation of the cervical facet joints in rats. *Spine (Phila Pa 1976)* 2001; 26:147-150.
35. Barnsley L, Bogduk N. Medial branch blocks are specific for the diagnosis of cervical zygapophyseal joint pain. *Reg Anesth* 1993; 18:343-350.
36. Zhang J, Suzuki N, Hirabayashi S, Saiki K, Fujita K. Surgical anatomy of the nerves and muscles in the posterior cervical spine. *Spine (Phila Pa 1976)* 2003; 28:1379-1384.
37. Siegenthaler A, Schliessbach J, Curatolo M, Eichenberger U. Ultrasound anatomy of the nerves supplying the cervical zygapophyseal joints: An exploratory study. *Reg Anesth Pain Med* 2011; 36:606-610.
38. Kallakuri S, Li Y, Chen C, Cavanaugh JM. Innervation of cervical ventral facet joint capsule: Histological evidence. *World J Orthop* 2012; 3:10-14.
39. Chua WH, Bogduk N. The surgical anatomy of thoracic facet denervation. *Acta Neurochir* 1995; 136:140-144.
40. Steinke H, Saito T, Miyaki T, Oi Y, Itoh M, Spanel-Borowski K. Anatomy of the human thoracolumbar Rami dorsales nervi spinalis. *Ann Anat* 2009; 191:408-416.
41. Sato T, Koizumi M, Kim JH, Kim JH, Wang BJ, Murakami G, Cho BH. Fetal development of deep back muscles in the human thoracic region with a focus on transversospinalis muscles and the medial branch of the spinal nerve posterior ramus. *J Anat* 2011; 219:756-765.
42. Marks RC. Distribution of pain provoked from lumbar facet joints and related structures during diagnostic spinal infiltration. *Pain* 1989; 39:37-40.
43. Fukui S, Ohseto K, Shiotani M, Ohno K, Karasawa H, Naganuma Y. Distribution of referred pain from the lumbar zygapophyseal joints and dorsal rami. *Clin J Pain* 1997; 13:303-307.
44. Hirsch C, Ingelmark BE, Miller M. The anatomical basis for low back pain. Studies on the presence of sensory nerve endings in ligamentous, capsular and intervertebral disc structures in the human lumbar spine. *Acta Orthop Scand* 1963; 33:1-17.
45. Windsor RE, King FJ, Roman SJ, Tata NS, Cone-Sullivan LA, Thmapi S, Acebey M, Gilhool JJ, Rao R, Sugar R. Electrical stimulation induced lumbar medial branch referral patterns. *Pain Physician* 2002; 5:347-353.
46. Mooney V, Robertson J. The facet syndrome. *Clin Orthop Relat Res* 1976; 115:149-156.
47. McCall IW, Park WM, O'Brien JP. Induced pain referral from posterior lumbar elements in normal subjects. *Spine (Phila Pa 1976)* 1979; 4:441-446.
48. Dreyfuss P, Tibiletti C, Dreyer SJ. Thoracic zygapophyseal joint pain patterns: A study in normal volunteers. *Spine (Phila Pa 1976)* 1994; 19:807-811.
49. Fukui S, Ohseto K, Shiotani M. Patterns of pain induced by distending the thoracic zygapophyseal joints. *Reg Anesth* 1997; 22:332-336.
50. Dwyer A, Aprill C, Bogduk N. Cervical zygapophyseal joint pain patterns: A study in normal volunteers. *Spine (Phila Pa 1976)* 1990; 15:453-457.
51. Aprill C, Dwyer A, Bogduk N. The prevalence of cervical zygapophyseal joint pain patterns II: A clinical evaluation. *Spine (Phila Pa 1976)* 1990; 15:458-461.
52. Fukui S, Ohseto K, Shiotani M, Ohno K, Karasawa H, Naganuma Y, Yuda Y. Referred pain distribution of the cervical zygapophyseal joints and cervical dorsal rami. *Pain* 1996; 68:79-83.
53. Windsor RE, Nagula D, Storm S. Electrical stimulation induced cervical medial branch referral patterns. *Pain Physician* 2003; 6:411-418.
54. Bogduk N. International Spinal Injection Society guidelines for the performance of spinal injection procedures. Part 1: Zygapophyseal joint blocks. *Clin J Pain* 1997; 13:285-302.
55. Falco FJE, Manchikanti L, Datta S, Sehgal N, Geffert S, Onyewu O, Zhu J, Coubarous S, Hameed M, Ward SP, Sharma M, Hameed H, Singh V, Boswell MV. An update of the effectiveness of therapeutic lumbar facet joint interventions. *Pain Physician* 2012; 15:E909-E953.
56. Falco FJE, Manchikanti L, Datta S, Wargo BW, Geffert S, Bryce DA, Atluri S, Singh V, Benyamin RM, Sehgal N, Ward S, Helm II S, Gupta S, Boswell MV. Systematic review of therapeutic effectiveness of cervical facet joint interventions: An update. *Pain Physician* 2012; 15:E839-E868.
57. Manchikanti KN, Atluri S, Singh V, Geffert S, Sehgal N, Falco FJE. An update of evaluation of therapeutic thoracic facet joint interventions. *Pain Physician* 2012; 15:E463-E481.
58. Leggett LE, Soril LJ, Lorenzetti DL, Noseworthy T, Steadman R, Tiwana S, Clement F. Radiofrequency ablation for chronic low back pain: A systematic review of randomized controlled trials. *Pain Res Manag* 2014; 19:e146-e153.
59. Poetscher AW, Gentil AF, Lenza M, Ferretti M. Radiofrequency denervation for facet joint low back pain: A systematic review. *Spine (Phila Pa 1976)* 2014; 39:E842-E849.
60. Bogduk N, Dreyfuss P, Govind J. A narrative review of lumbar medial branch neurotomy for the treatment of back pain. *Pain Med* 2009; 10:1035-1045.
61. Bogduk N. A narrative review of intraarticular corticosteroid injections for low back pain. *Pain Med* 2005; 6:287-296.
62. Manchikanti L, Datta S, Gupta S, Munglani R, Bryce DA, Ward SP, Benyamin RM, Sharma ML, Helm II S, Fellows B, Hirsch JA. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: Part 2. Therapeutic interventions. *Pain Physician* 2010; 13:E215-E264.
63. Manchikanti L, Singh V, Derby R, Schultz DM, Benyamin RM, Prager JP, Hirsch JA. Reassessment of evidence synthesis of occupational medicine practice guidelines for interventional pain management. *Pain Physician* 2008; 11:393-482.
64. Niemisto L, Kalso E, Malmivaara A, Seit-salo S, Hurri H. Radiofrequency denervation for neck and back pain. A systematic review of randomized controlled trials. *Cochrane Database Syst Rev* 2003; CD004058.
65. Manchikanti L, Benyamin RM, Falco FJE, Caraway DL, Datta S, Hirsch JA. Guidelines warfare over interventional techniques: Is there a lack of discourse or straw man? *Pain Physician* 2012; 15:E1-E26.
66. Chou R, Hashimoto R, Friedly J, Fu Rochelle, Dana T, Sullivan S, Bougatsos C, Jarvik J. Pain Management Injection Therapies for Low Back Pain. Technology Assessment Report ESIB0813. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. HHS 290-2012-00014-I.) Rockville, MD: Agency for Healthcare Research and Quality; March 20, 2015. [www.cms.gov/Medicare/Coverage/DeterminationProcess/Downloads/id98TA.pdf](http://www.cms.gov/Medicare/Coverage/DeterminationProcess/Downloads/id98TA.pdf)
67. Health Technology Assessment, Washington State Health Care Authority. Spi-

- nal Injections, Updated Final Evidence Report. Spectrum Research, Inc., March 10, 2011.
68. Database of Abstracts of Reviews of Effects (DARE): Quality-assessed Reviews. York (UK): Centre for Reviews and Dissemination. An update of the effectiveness of therapeutic lumbar facet joint interventions. 2012 (cited 2013 Feb 21).
  69. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *Ann Intern Med* 2009; 151:W65-W94.
  70. Furlan AD, Pennick V, Bombardier C, van Tulder M; Editorial Board, Cochrane Back Review Group. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine (Phila Pa 1976)* 2009; 34:1929-1941.
  71. Manchikanti L, Falco FJE, Singh V, Benyamin RM, Racz GB, Helm II S, Caraway DL, Calodney AK, Snook LT, Smith HS, Gupta S, Ward SP, Grider JS, Hirsch JA. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain. Part I: Introduction and general considerations. *Pain Physician* 2013; 16:S1-S48.
  72. Manchikanti L, Hirsch JA, Cohen SP, Heavner JE, Falco FJE, Diwan S, Boswell MV, Candido KD, Onyewu O, Zhu J, Sehgal N, Kaye AD, Benyamin RM, Helm II S, Singh V, Datta S, Abdi S, Christo PJ, Hameed H, Hameed M, Vallejo R, Pampati V, Racz GB, Raj PP. Assessment of methodologic quality of randomized trials of interventional techniques: Development of an interventional pain management specific instrument. *Pain Physician* 2014; 17:E263-E290.
  73. Manchikanti L, Hirsch JA, Heavner JE, Cohen SP, Benyamin RM, Sehgal N, Falco FJE, Vallejo R, Onyewu O, Zhu J, Kaye AD, Boswell MV, Helm II S, Candido KD, Diwan S, Simopoulos TT, Singh V, Pampati V, Racz GB, Raj PP. Development of an interventional pain management specific instrument for methodologic quality assessment of nonrandomized studies of interventional techniques. *Pain Physician* 2014; 17:E291-E317.
  74. Farrar JT. What is clinically meaningful: Outcome measures in pain clinical trials. *Clin J Pain* 2000; 16:S106-S112.
  75. Salaffi F, Stancati A, Silvestri CA, Ciapetti A, Grassi W. Minimal clinically important changes in chronic musculoskeletal pain intensity measured on a numerical rating scale. *Eur J Pain* 2004; 8:283-291.
  76. Hagg O, Fritzell P, Nordwall A. The clinical importance of changes in outcome scores after treatment for chronic low back pain. *Eur Spine J* 2003; 12:12-20.
  77. Carragee EJ. The rise and fall of the "minimum clinically important difference." *Spine J* 2010; 10:283-284.
  78. Gatchel RJ, Mayer TG, Choi Y, Chou R. Validation of a consensus-based minimal clinically important difference (MCID) threshold using an objective functional external anchor. *Spine J* 2013; 13:889-893.
  79. Parker SL, Adogwa O, Mendenhall SK, Shau DN, Anderson WN, Cheng JS, Devin CJ, McGirt MJ. Determination of minimum clinically important difference (MCID) in pain, disability, and quality of life after revision fusion for symptomatic pseudoarthrosis. *Spine J* 2012; 12:1122-1128.
  80. Copay AG, Subach BR, Glassman SD, Polly DW Jr, Schuler TC. Understanding the minimum clinically important difference: A review of concepts and methods. *Spine J* 2007; 7:541-546.
  81. Rouquette A, Blanchin M, Sébille V, Guillemin F, Côté SM, Falissard B, Harouin JB. The minimal clinically important difference determined using item response theory models: An attempt to solve the issue of the association with baseline score. *J Clin Epidemiol* 2014; 67:433-440.
  82. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V. Evaluation of lumbar facet joint nerve blocks in managing chronic low back pain: A randomized, double-blind, controlled trial with a 2-year follow-up. *Int J Med Sci* 2010; 7:124-135.
  83. Manchikanti L, Singh V, Falco FJE, Cash KA, Fellows B. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: A randomized, double-blind controlled trial. *Pain Physician* 2010; 13:437-450.
  84. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V, Fellows B. The role of thoracic medial branch blocks in managing chronic mid and upper back pain: A randomized, double-blind, active-control trial with a 2-year follow-up. *Anesthesiol Res Pract* 2012; 2012:585806.
  85. Manchikanti L, Cash KA, McManus CD, Pampati V. Assessment of effectiveness of percutaneous adhesiolysis in managing chronic low back pain secondary to lumbar central spinal canal stenosis. *Int J Med Sci* 2013; 10:50-59.
  86. Manchikanti L, Singh V, Cash KA, Pampati V, Datta S. Assessment of effectiveness of percutaneous adhesiolysis and caudal epidural injections in managing lumbar post surgery syndrome: A 2-year follow-up of randomized, controlled trial. *J Pain Res* 2012; 5:597-608.
  87. Manchikanti L, Singh V, Cash KA, Pampati V, Damron KS, Boswell MV. Effect of fluoroscopically guided caudal epidural steroid or local anesthetic injections in the treatment of lumbar disc herniation and radiculitis: A randomized, controlled, double blind trial with a two-year follow-up. *Pain Physician* 2012; 15:273-286.
  88. Manchikanti L, Singh V, Cash KA, Pampati V, Datta S. Fluoroscopic caudal epidural injections in managing post lumbar surgery syndrome: Two-year results of a randomized, double-blind, active-control trial. *Int J Med Sci* 2012; 9:582-591.
  89. Manchikanti L, Cash KA, McManus CD, Pampati V, Fellows B. Results of 2-year follow-up of a randomized, double-blind, controlled trial of fluoroscopic caudal epidural injections in central spinal stenosis. *Pain Physician* 2012; 15:371-384.
  90. Manchikanti L, Singh V, Cash KA, Pampati V, Falco FJE. A randomized, double-blind, active-control trial of the effectiveness of lumbar interlaminar epidural injections in disc herniation. *Pain Physician* 2014; 17:E61-E74.
  91. Manchikanti L, Cash KA, McManus CD, Pampati V, Benyamin RM. A randomized, double-blind, active-controlled trial of fluoroscopic lumbar interlaminar epidural injections in chronic axial or discogenic low back pain: Results of a 2-year follow-up. *Pain Physician* 2013; 16:E491-E504.
  92. Manchikanti L, Cash KA, McManus CD, Damron KS, Pampati V, Falco FJE. A randomized, double-blind controlled trial of lumbar interlaminar epidural injections in central spinal stenosis: 2-year follow-up. *Pain Physician* 2015; 18:79-92.
  93. Manchikanti L, Cash KA, Pampati V, Falco FJE. Transforaminal epidural injections in chronic lumbar disc herniation: A randomized, double-blind, active-control trial. *Pain Physician* 2014; 17:E489-E501.
  94. Manchikanti L, Cash KA, Pampati V, Malla Y. Two-year follow-up results of fluoroscopic cervical epidural injections

- in chronic axial or discogenic neck pain: A randomized, double-blind, controlled trial. *Int J Med Sci* 2014; 11:309-320.
95. Manchikanti L, Cash KA, Pampati V, Wargo BW, Malla Y. A randomized, double-blind, active control trial of fluoroscopic cervical interlaminar epidural injections in chronic pain of cervical disc herniation: Results of a 2-year follow-up. *Pain Physician* 2013; 16:465-478.
  96. Manchikanti L, Malla Y, Cash KA, McManus CD, Pampati V. Fluoroscopic epidural injections in cervical spinal stenosis: Preliminary results of a randomized, double-blind, active control trial. *Pain Physician* 2012; 15:E59-E70.
  97. Manchikanti L, Malla Y, Cash KA, McManus CD, Pampati V. Fluoroscopic cervical interlaminar epidural injections in managing chronic pain of cervical post-surgery syndrome: Preliminary results of a randomized, double-blind active control trial. *Pain Physician* 2012; 15:13-26.
  98. Lohr KN, Carey TS. Assessing 'best evidence': Issues in grading the quality of studies for systematic reviews. *J Comm J Qual Improv* 1999; 25:470-479.
  99. Clarke M, Oxman AD. Cochrane Reviewer's Handbook 4.0. The Cochrane Collaboration, 1999.
  100. National Health and Medical Research Council (NHMRC). How to Use the Evidence: Assessment and Application of Scientific Evidence. NHMRC, Canberra, Australia, 2000.
  101. Staal JB, de Bie RA, de Vet HC, Hildebrandt J, Nelemans P. Injection therapy for subacute and chronic low back pain: An updated Cochrane review. *Spine (Phila Pa 1976)* 2009; 34:49-59.
  102. Harris RP, Helfand M, Woolf SH, Lohr KN, Mulrow CD, Teutsch SM, Atkins D; Methods Work Group, Third US Preventive Services Task Force. Current methods of the US Preventive Services Task Force. *Am J Prevent Med* 2001; 20:21-35.
  103. Guyatt GH, Cook DJ, Sackett DL, Eckman M, Pauker S. Grades of recommendation for antithrombotic agents. *Chest* 1998; 114:441S-444S.
  104. Gross PA, Barrett TL, Dellinger EP, Krause PJ, Martone WJ, McGowan JE Jr, Sweet RL, Wenzel RP. Purpose of quality standards for infectious diseases. Infectious Diseases Society of America. *Clin Infect Dis* 1994; 18:421.
  105. Manchikanti L, Falco FJE, Benjamin RM, Kaye AD, Boswell MV, Hirsch JA. A modified approach to grading of evidence. *Pain Physician* 2014; 17:E319-E325.
  106. Jeong SY, Kim JS, Choi WS, Hur JW, Ryu KS. The effectiveness of endoscopic radiofrequency denervation of medial branch for treatment of chronic low back pain. *J Korean Neurosurg Soc* 2014; 56:338-343.
  107. Li ZZ, Hou SX, Shang WL, Song KR, Wu WW. Evaluation of endoscopic dorsal ramus rhizotomy in managing facetogenic chronic low back pain. *Clin Neurol Neurosurg* 2014; 126:11-17.
  108. Cohen SP, Williams KA, Kurihara C, Nguyen C, Shields C, Kim P, Griffith SR, Larkin TM, Crooks M, Williams N, Morlando B, Strassels SA. Multicenter, randomized, comparative cost-effectiveness study comparing 0, 1, and 2 diagnostic medial branch (facet joint nerve) block treatment paradigms before lumbar facet radiofrequency denervation. *Anesthesiology* 2010; 113:395-405.
  109. Manchikanti L, Singh V, Falco FJ, Cash KA, Pampati V. Lumbar facet joint nerve blocks in managing chronic facet joint pain: One-year follow-up of a randomized, double-blind controlled trial: Clinical Trial NCT00355914. *Pain Physician* 2008; 11:121-132.
  110. Carette S, Marcoux S, Truchon R, Grondin C, Gagnon J, Allard Y, Latulippe M. A controlled trial of corticosteroid injections into facet joints for chronic low back pain. *N Engl J Med* 1991; 325:1002-1007.
  111. Fuchs S, Erbe T, Fischer HL, Tibesku CO. Intraarticular hyaluronic acid versus glucocorticoid injections for nonradicular pain in the lumbar spine. *J Vasc Interv Radiol* 2005; 16:1493-1498.
  112. Manchikanti L, Pampati V, Bakht C, Rivera J, Beyer C, Damron K, Barnhill R. Effectiveness of lumbar facet joint nerve blocks in chronic low back pain: A randomized clinical trial. *Pain Physician* 2001; 4:101-117.
  113. Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: A randomized double-blind trial. *Spine (Phila Pa 1976)* 2008; 33:1291-1298.
  114. van Wijk RM, Geurts JW, Wynne HJ, Hammink E, Buskens E, Lousberg R, Knape JT, Groen GJ. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: A randomized, double-blind, sham lesion-controlled trial. *Clin J Pain* 2005; 21:335-344.
  115. van Kleef M, Barendse GAM, Kessels A, Voets HM, Weber WE, de Lange S. Randomized trial of radiofrequency lumbar facet denervation for chronic low back pain. *Spine (Phila Pa 1976)* 1999; 24:1937-1942.
  116. Tekin I, Mirzai H, Ok G, Erbuyun K, Vatansever D. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain* 2007; 23:524-529.
  117. Civelek E, Cansever T, Kabatas S, Kircelli A, Yilmaz C, Muslumman M, Ofluoglu D, Caner H. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg* 2012; 22:200-206.
  118. Dobrogowski J, Wrzosek A, Wordliczek J. Radiofrequency denervation with or without addition of pentoxifylline or methylprednisolone for chronic lumbar zygapophysial joint pain. *Pharmacol Rep* 2005; 57:475-480.
  119. Barnsley L, Lord SM, Wallis BJ, Bogduk N. Lack of effect of intra-articular corticosteroids for chronic pain in the cervical zygapophysial joints. *N Engl J Med* 1994; 330:1047-1050.
  120. Joo YC, Park JY, Kim KH. Comparison of alcohol ablation with repeated thermal radiofrequency ablation in medial branch neurotomy for the treatment of recurrent thoracolumbar facet joint pain. *J Anesth* 2013; 27:390-395.
  121. Ribeiro LH, Furtado RN, Konai MS, Andreo AB, Rosenfeld A, Natour J. Effect of facet joint injection versus systemic steroids in low back pain: A randomized controlled trial. *Spine (Phila Pa 1976)* 2013; 38:1995-2002.
  122. Moon JY, Lee PB, Kim YC, Choi SP, Sim WS. An alternative distal approach for the lumbar medial branch radiofrequency denervation: A prospective randomized comparative study. *Anesth Analg* 2013; 116:1133-1140.
  123. Lakemeier S, Lind M, Schultz W, Fuchs-Winkelmann S, Timmesfeld N, Foelsch C, Peterlein CD. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: A randomized, controlled, double-blind trial. *Anesth Analg* 2013; 117:228-235.
  124. Yun DH, Kim HS, Yoo SD, Kim DH, Chon JM, Choi SH, Hwang DG, Jung PK. Efficacy of ultrasonography-guided injections in patients with facet syndrome of the low lumbar spine. *Ann Rehabil Med* 2012; 36:66-71.

125. Obernauer J, Galiano K, Gruber H, Bale R, Obwegeser AA, Schatzer R, Loizides A. Ultrasound-guided versus computed tomography-controlled facet joint injections in the middle and lower cervical spine: A prospective randomized clinical trial. *Med Ultrason* 2013; 15:10-15.
126. Wen CB, Li YZ, Sun L, Xiao H, Yang BX, Song L, Liu H. A clinical trial of ultrasound-guided facet joint block in the lumbar spine to treat facet joint related low back pain. *Sichuan Da Xue Xue Bao Yi Xue Ban* 2014; 45:712-716.
127. Klessinger S. Zygapophysial joint pain in post lumbar surgery syndrome. The efficacy of medial branch blocks and radiofrequency neurotomy. *Pain Med* 2013; 14:374-377.
128. Lord S, Barnsley L, Wallis B, McDonald G, Bogduk N. Percutaneous radio-frequency neurotomy for chronic cervical zygapophyseal-joint pain. *N Engl J Med* 1996; 335:1721-1726.
129. Park SC, Kim KH. Effect of adding cervical facet joint injections in a multimodal treatment program for long-standing cervical myofascial pain syndrome with referral pain patterns of cervical facet joint syndrome. *J Anesth* 2012; 26:738-745.
130. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V, Fellows B. Comparative effectiveness of a one-year follow-up of thoracic medial branch blocks in management of chronic thoracic pain: A randomized, double-blind active controlled trial. *Pain Physician* 2010; 13:535-548.
131. Lilius G, Laasonen EM, Myllynen P, Harilainen A, Gronlund G. Lumbar facet joint syndrome. A randomised clinical trial. *J Bone Joint Surg Br* 1989; 71:681-684.
132. Marks RC, Houston T, Thulbourne T. Facet joint injection and facet nerve block: A randomised comparison in 86 patients with chronic low back pain. *Pain* 1992; 49:325-328.
133. Nash TP. Facet joints. Intra-articular steroids or nerve blocks? *Pain Clinic* 1990; 3:77-82.
134. Leclaire R, Fortin L, Lambert R, Bergeron YM, Rossignol M. Radiofrequency facet joint denervation in the treatment of low back pain: A placebo-controlled clinical trial to assess efficacy. *Spine (Phila Pa 1976)* 2001; 26:1411-1416.
135. Gallagher J, Petriccione Di Vadi PL, Wedley JR, Hamann W, Ryan P, Chikanza I, Kirkham B, Price R, Watson MS, Grahme R, Wood S. Radiofrequency facet joint denervation in the treatment of low back pain: A prospective controlled double-blind study to assess its efficacy. *Pain Clinic* 1994; 7:193-198.
136. Kroll H, Kim D, Danic M, Sankey S, Gariwala M, Brown M. A randomized, double-blind, prospective study comparing the efficacy of continuous versus pulsed radiofrequency in the treatment of lumbar facet syndrome. *J Clin Anesth* 2008; 20:534-537.
137. Ackerman WE 3rd, Ahmad M. Pain relief with intraarticular or medial branch nerve blocks in patients with positive lumbar facet joint SPECT imaging: A 12-week outcome study. *South Med J* 2008; 101:931-934.
138. Andres RH, Graupner T, Bärlocher CB, Augsburger A, Fandino J. Laser-guided lumbar medial branch cryorhizotomy. *J Neurosurg Spine* 2010; 13:341-345.
139. MacVicar J, Borowczyk JM, MacVicar AM, Loughnan BM, Bogduk N. Lumbar medial branch radiofrequency neurotomy in New Zealand. *Pain Med* 2013; 14:639-645.
140. Kader D, Radha S, Smith F, Wardlaw D, Scott N, Rege A, Pope M. Evaluation of perifacet injections and paraspinal muscle rehabilitation in treatment of low back pain. A randomised controlled trial. *Ortop Traumatol Rehabil* 2012; 14:251-259.
141. Tzaan W, Tasker R. Percutaneous radiofrequency facet rhizotomy—experience with 118 procedures and reappraisal of its value. *Can J Neurol Sci* 2000; 27:125-130.
142. Stolker RJ, Vervest AC, Groen GJ. Parameters in electrode positioning in thoracic percutaneous facet denervation: An anatomical study. *Acta Neurochir* 1994; 128:32-39.
143. Stolker RJ, Vervest AC, Groen GJ. The treatment of chronic thoracic segmental pain by radiofrequency percutaneous partial rhizotomy. *J Neurosurg* 1994; 80:986-992.
144. Chua NH, Vissers KC, Sluijter ME. Pulsed radiofrequency treatment in interventional pain management: Mechanisms and potential indications – a review. *Acta Neurochir (Wien)* 2011; 153:763-771.
145. Golovac S. Radiofrequency neurolysis. *Neuroimaging Clin N Am* 2010; 20:203-214.
146. Mitra R, Do H, Alamin T, Cheng I. Facet pain in thoracic compression fractures. *Pain Med* 2010; 11:1674-1677.
147. Haufe SM, Mork AR. Endoscopic facet debridement for the treatment of facet arthritic —pain — A novel new technique. *Int J Med Sci* 2010; 7:120-123.
148. Slappendel R, Crul BJ, Braak GJ, Geurts JW, Booi LH, Voerman VF, de Boo T. The efficacy of radiofrequency lesioning of the cervical spinal dorsal root ganglion in a double blinded randomized study: No difference between 40°C and 67°C treatments. *Pain* 1997; 73:159-163.
149. Haspelslagh SR, Van Suijlekom HA, Lame IE, Kessels AG, van Kleef M, Weber WE. Randomised controlled trial of cervical radiofrequency lesions as a treatment for cervicogenic headache: ISRCTN07444684. *BMC Anesthesiol* 2006; 6:1.
150. Wallis B, Lord S, Bogduk N. Resolution of psychological distress of whiplash patients following treatment by radiofrequency neurotomy: A randomized, double-blind, placebo-controlled trial. *Pain* 1997; 73:15-22.
151. Schaerer JP. Radiofrequency facet rhizotomy in the treatment of chronic neck and low back pain. *Int Surg* 1978; 63:53-59.
152. Manchikanti L, Manchikanti K, Damron K, Pampati V. Effectiveness of cervical medial branch blocks in chronic neck pain: A prospective outcome study. *Pain Physician* 2004; 7:195-201.
153. Roy C, Chatterjee N, Ganguly S, Sengupta R. Efficacy of combined treatment with medial branch radiofrequency neurotomy and steroid block in lumbar facet joint arthropathy. *J Vasc Interv Radiol* 2012; 23:1659-1664.
154. Mehnert MJ, Freedman MK. Update on the role of z-joint injection and radiofrequency neurotomy for cervicogenic headache. *PM R* 2013; 5:221-227.
155. Kim D. Bipolar intra-articular radiofrequency thermocoagulation of the thoracic facet joints: A case series of a new technique. *Korean J Pain* 2014; 27:43-48.
156. Saltychev M, Laimi K. Re: Poetscher AW, Gentil AF, Lenza M, et al. Radiofrequency denervation for facet joint low back pain: A systematic review. *Spine (Phila Pa 1976)* 2014; 39:E842-9. *Spine (Phila Pa 1976)* 2014; 39:E1498-E1500.
157. Carragee EJ, Hurwitz EL, Cheng I, Carroll LJ, Nordin M, Guzman J, Peloso P, Holm LW, Côté P, Hogg-Johnson S, van der Velde G, Cassidy JD, Haldeman S; Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. Treatment of neck pain: In-

- jections and surgical interventions: Results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)* 2008; 33:S153-S169.
158. Dreyfuss P, Baker R. In response to treatment of neck pain. *Eur Spine J* 2008; 17:1270-1272; author reply 1273-1275.
  159. Sapir DA, Gorup JM. Radiofrequency medial branch neurotomy in litigant and non-litigant patients with cervical whiplash. *Spine (Phila Pa 1976)* 2001; 26:E268-E273.
  160. MacVicar J, Borowczyk JM, MacVicar AM, Loughnan BM, Bogduk N. Cervical medial branch radiofrequency neurotomy in New Zealand. *Pain Med* 2012; 13:647-654.
  161. Speldewinde GC. Outcomes of percutaneous zygapophysial and sacroiliac joint neurotomy in a community setting. *Pain Med* 2011; 12:209-218.
  162. Manchikanti L, Manchikanti KN, Manchukonda R, Pampati V, Cash KA. Evaluation of therapeutic thoracic medial branch block effectiveness in chronic thoracic pain: A prospective outcome study with minimum 1-year follow up. *Pain Physician* 2006; 9:97-105.
  163. McDonald G, Lord S, Bogduk N. Long-term follow-up of patients treated with cervical radiofrequency neurotomy for chronic spinal pain. *Neurosurgery* 1999; 45:61-67.
  164. Barnsley L. Percutaneous radiofrequency neurotomy for chronic neck pain: Outcomes in a series of consecutive patients. *Pain Med* 2005; 6:282-286.
  165. Leclaire R, Fortin L, Bergeron Y, Lambert R, Rissignol M. In response to: Radiofrequency neurotomy for chronic lumbar facet pain – interpreting the evidence. *Pain Pract* 2010; 10:261-262.
  166. Manchikanti L, Pampati V, Damron KS. The role of placebo and nocebo effects of perioperative administration of sedatives and opioids in interventional pain management. *Pain Physician* 2005; 8:349-355.
  167. Manchikanti L, Giordano J, Fellows B, Hirsch JA. Placebo and nocebo in interventional pain management: A friend or a foe – or simply foes? *Pain Physician* 2011; 14:E157-E175.
  168. Wells RE, Kaptchuk TJ. To tell the truth, the whole truth, may do patients harm: The problem of the nocebo effect for informed consent. *Am J Bioeth* 2012; 12:22-29.
  169. Häuser W, Bartram C, Bartram-Wunn E, Tölle T. Adverse events attributable to nocebo in randomized controlled drug trials in fibromyalgia syndrome and painful diabetic peripheral neuropathy: Systematic review. *Clin J Pain* 2012; 28:437-451.
  170. Lyby PS, Forsberg JT, Asli O, Flaten MA. Induced fear reduces the effectiveness of a placebo intervention on pain. *Pain* 2012; 153:1114-1121.
  171. Hróbjartsson A, Gøtzsche PC. Placebo interventions for all clinical conditions. *Cochrane Database Syst Rev* 2010; 1:CD003974.
  172. Bingel U; Placebo Competence Team. Avoiding nocebo effects to optimize treatment outcome. *JAMA* 2014; 312:693-694.
  173. Colloca L, Finniss D. Nocebo effects, patient-clinician communication, and therapeutic outcomes. *JAMA* 2012; 307:567-568.
  174. Howick J, Bishop FL, Heneghan, Wolstenholme J, Stevens S, Hobbs FDR, Lewith G. Placebo use in the United Kingdom: Results from a national survey of primary care practitioners. *PLoS One* 2013; 8:e8247.
  175. Howick J, Friedemann C, Tsakok M, Watson R, Tsakok T, Thomas J, Perera R, Fleming S, Heneghan C. Are treatments more effective than placebos? A systematic review and meta-analysis. *PLoS One* 2013; 8:e62599.
  176. Kaptchuk TJ, Friedlander E, Kelley JM, Sanchez MN, Kokkotou E, Singer JP, Kowalczykowski M, Miller FG, Kirsch I, Lembo AJ. Placebos without deception: A Randomized controlled trial in irritable bowel syndrome. *PLoS One* 2010; 5: e15591.
  177. Petersen GL, Finnerup NB, Colloca L, Amanzio M, Price DD, Jensen TS, Vase L. The magnitude of nocebo effects in pain: A meta-analysis. *Pain* 2014; 155:1426-1434.
  178. Ghai B, Kumar K, Bansal D, Dhatt SS, Kanukula R, Batra YK. Effectiveness of parasagittal interlaminar epidural local anesthetic with or without steroid in chronic lumbosacral pain: A randomized, double-blind clinical trial. *Pain Physician* 2015; 18:237-248.
  179. Ghai B, Bansal D, Kay JP, Vadaje KS, Wig J. Transforaminal versus parasagittal interlaminar epidural steroid injection in low back pain with radicular pain: A randomized, double-blind, active-control trial. *Pain Physician* 2015; 17:277-290.
  180. Manchikanti L, Nampiaparampil DE, Manchikanti KN, Falco FJE, Singh V, Benyamin RM, Kaye AD, Sehgal N, Soin A, Simopoulos TT, Bakshi S, Gharibo CG, Gilligan CJ, Hirsch JA. Comparison of the efficacy of saline, local anesthetics, and steroids in epidural and facet joint injections for the management of spinal pain: A systematic review of randomized controlled trials. *Surg Neurol Int* 2015; 6:S194-S235.
  181. Nagar VR, Birthi P, Grider JS, Asopa A. Systematic review of radiofrequency ablation and pulsed radiofrequency for management of cervicogenic headache. *Pain Physician* 2015; 18:109-130.
  182. Manchikanti L, Kaye AD, Manchikanti KN, Boswell MV, Pampati V, Hirsch JA. Efficacy of epidural injections in the treatment of lumbar central spinal stenosis: A systematic review. *Anesth Pain Med* 2015; 5:e23139.
  183. Manchikanti L, Benyamin RM, Falco FJ, Kaye AD, Hirsch JA. Do epidural injections provide short- and long-term relief for lumbar disc herniation? A systematic review. *Clin Orthop Relat Res* 2015; 473:1940-1956.
  184. Manchikanti L, Nampiaparampil DE, Candido KD, Bakshi S, Grider JS, Falco FJE, Sehgal N, Hirsch JA. Do cervical epidural injections provide long-term relief in neck and upper extremity pain? A systematic review. *Pain Physician* 2015; 18:39-60.
  185. Lee HM, Weinstein JN, Meller ST, Hayashi N, Spratt KF, Gebhart GF. The role of steroids and their effects on phospholipase A2. An animal model of radiculopathy. *Spine (Phila Pa 1976)* 1998; 23:1191-1196.
  186. Johansson A, Hao J, Sjölund B. Local corticosteroid application blocks transmission in normal nociceptive C-fibres. *Acta Anaesthesiol Scand* 1990; 34:335-338.
  187. Pasqualucci A, Varrassi G, Braschi A, Peduto VA, Brunelli A, Marinangeli F, Gori F, Colò F, Paladín A, Mojoli F. Epidural local anesthetic plus corticosteroid for the treatment of cervical brachial radicular pain: Single injection versus continuous infusion. *Clin J Pain* 2007; 23:551-557.
  188. Sato C, Sakai A, Ikeda Y, Suzuki H, Sakamoto A. The prolonged analgesic effect of epidural ropivacaine in a rat model of neuropathic pain. *Anesth Analg* 2008; 106:313-320.
  189. Tachihara H, Sekiguchi M, Kikuchi S, Konno S. Do corticosteroids produce additional benefit in nerve root infiltration for lumbar disc herniation. *Spine (Phila Pa 1976)* 2008; 33:743-747.

190. Arner S, Lindblom U, Meyerson BA, Molander C. Prolonged relief of neuralgia after regional anesthetic block. A call for further experimental and systematic clinical studies. *Pain* 1990; 43:287-297.
191. Lavoie PA, Khazen T, Filion PR. Mechanisms of the inhibition of fast axonal transport by local anesthetics. *Neuropharmacology* 1989; 28:175-181.
192. Bisby MA. Inhibition of axonal transport in nerves chronically treated with local anesthetics. *Exp Neurol* 1975; 47:481-489.
193. Katz WA, Rothenberg R. The nature of pain: Pathophysiology. *J Clin Rheumatol* 2005; 11:511-515.
194. Melzack R,Coderre TJ, Katz J, Vaccarino AL. Central neuroplasticity and pathological pain. *Ann N Y Acad Sci* 2001; 933:157-174.
195. Cassuto J, Sinclair R, Bonderovic M. Anti-inflammatory properties of local anesthetics and their present and potential clinical implications. *Acta Anaesthesiol Scand* 2006; 50:265-282.
196. Carette S, Leclaire R, Marcoux S, Morin F, Blaise GA, St-Pierre A, Truchon R, Parent F, Levesque J, Bergeron V, Montminy P, Blanchette C. Epidural corticosteroid injections for sciatica due to herniated nucleus pulposus. *N Engl J Med* 1997; 336:1634-1640.
197. Karppinen J, Malmivaara A, Kurunlahti M, Kyllönen E, Pienimäki T, Nieminen P, Ohinmaa A, Tervonen O, Vanharanta H. Periradicular infiltration for sciatica: A randomized controlled trial. *Spine (Phila Pa 1976)* 2001; 26:1059-1067.
198. Hayashi N, Weinstein JN, Meller ST, Lee HM, Spratt KF, Gebhart GF. The effect of epidural injection of betamethasone or bupivacaine in a rat model of lumbar radiculopathy. *Spine (Phila Pa 1976)* 1998; 23:877-885.
199. Manchikanti L, Datta S, Derby R, Wolfer LR, Benyamin RM, Hirsch JA. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: Part 1. Diagnostic interventions. *Pain Physician* 2010; 13:E141-E174.
200. Manchikanti L, Shah RV, Datta S, Singh V. Critical evaluation of interventional pain management literature provides inaccurate conclusions. *Spine J* 2009; 9:706-708.
201. Smuck M, Levin JH: RE: Manchikanti L, Singh V, Falco FJE, Cash KA, Fellows B. Cervical medial branch blocks for chronic cervical facet joint pain: A randomized double-blind, controlled trial with one-year follow-up. *Spine (Phila Pa 1976)* 2008; 33:1813-20. *Spine (Phila Pa 1976)* 2009; 34:1116.
202. Chou R, Atlas SJ, Stanos SP, Rosenquist RW. Nonsurgical interventional therapies for low back pain: A review of the evidence for an American Pain Society clinical practice guideline. *Spine (Phila Pa 1976)* 2009; 34:1078-1093.
203. Manchikanti L, Falco FJE, Boswell MV, Hirsch JA. Facts, fallacies, and politics of comparative effectiveness research: Part 2. Implications for interventional pain management. *Pain Physician* 2010; 13:E55-E79.
204. Chou R, Atlas SJ, Loeser JD, Rosenquist RW, Stanos SP. Guideline warfare over interventional therapies for low back pain: Can we raise the level of discourse? *J Pain* 2011; 12:833-839.
205. Birkenmaier C, Veihelmann A, Trouillier HH, Hausdorf J, von Schulze P, Pöhlgen C. Medial branch blocks versus pericapsular blocks in selecting patients for percutaneous cryodenervation of lumbar facet joints. *Reg Anesth Pain Med* 2007; 32:27-33.
206. Pham Dang C, Lelong A, Guilley J, Nguyen JM, Volteau C, Venet G, Perrier C, Lejus C, Blanloeil Y. Effect on neurostimulation of injectates used for perineural space expansion before placement of a stimulating catheter: Normal saline versus dextrose 5% in water. *Reg Anesth Pain Med* 2009; 34:398-403.
207. Indahl A, Kaigle AM, Reikerås O, Holm SH. Interaction between the porcine lumbar intervertebral disc, zygapophysial joints, and paraspinal muscles. *Spine (Phila Pa 1976)* 1997; 22:2834-2840.
208. Indahl A, Kaigle A, Reikerås O, Holm S. Electromyographic response of the porcine multifidus musculature after nerve stimulation. *Spine (Phila Pa 1976)* 1995; 20:2652-2658.
209. Johnson CR, Barr RC, Klein SM. A computer model of electrical stimulation of peripheral nerves in regional anesthesia. *Anesthesiology* 2007; 106:323-330.
210. Eippert F, Finsterbusch J, Bingel U, Büchel C. Direct evidence for spinal cord involvement in placebo analgesia. *Science* 2009; 326:404.
211. Tsui BC, Kropelin B, Ganapathy S, Finucane B. Dextrose 5% in water: Fluid medium maintaining electrical stimulation of peripheral nerve during stimulating catheter placement. *Acta Anaesthesiol Scand* 2005; 49:1562-1565.
212. Holm I, Friis A, Brox JI, Gunderson R, Steen H. Minimal influence of facet joint anesthesia on isokinetic muscle performance in patients with chronic degenerative low back disorders. *Spine (Phila Pa 1976)* 2000; 25:2091-2094.
213. Kang YM, Choi WS, Pickar JG. Electrophysiologic evidence for an intersegmental reflex pathway between lumbar paraspinal tissues. *Spine (Phila Pa 1976)* 2002; 27:E56-E63.
214. Bhatia MT, Parikh LCJ. Epidural saline therapy in lumbo-sciatic syndrome. *J Indian Med Assoc* 1966; 47:537-542.
215. Gupta AK, Mital VK, Azmi RU. Observations of the management of lumbosciatic syndromes (sciatica) by epidural saline. *J Indian Med Assoc* 1970; 54:194-196.
216. Wittenberg RH, Greskötter KR, Steffen R, Schoenfeld BL. Is epidural injection treatment with hypertonic saline solution in intervertebral disk displacement useful? (The effect of NaCl solution on intervertebral disk tissue). *Z Orthop Ihre Grenzgeb* 1990; 128:223-226.
217. Levin JH. Prospective, double-blind, randomized placebo-controlled trials in interventional spine: What the highest quality literature tells us. *Spine J* 2009; 9:690-703.
218. O'Neill C. Re: Manchikanti L, Singh V, Falco FJE, et al. Cervical medial branch blocks for chronic cervical facet joint pain. A randomized, double-blind, controlled trial with one-year follow-up. *Spine* 2008; 33:1813-20. *Spine (Phila Pa 1976)* 2009; 34:1117-1118.
219. Louhiala P, Puustinen R. Rethinking the placebo effect. *J Med Ethics* 2008; 34:107-109.
220. Louhiala P. The ethics of the placebo in clinical practice revisited. *J Med Ethics* 2009; 35:407-409.
221. Concato J. When to randomize, or "evidence-based medicine needs medicine-based evidence." *Pharmacoepidemiol Drug Saf* 2012; 21:6-12.
222. Quill TE, Holloway RG. Evidence, preferences, —recommendations — finding the right balance in patient care. *N Engl J Med* 2012; 366:1653-1655.
223. Gelijns AC, Gabriel SE. Looking beyond translation — integrating clinical research with medical practice. *N Engl J Med* 2012; 366:1659-1661.
224. Blease C. The principle of parity: The "placebo effect" and physician communication. *J Med Ethics* 2012; 38:199-203.
225. Manchikanti L, Benyamin RM. Key safety considerations when administering epidural steroid injections. *Pain Manag* 2015; Jun 10:1-12. [Epub ahead of print]
226. Manchikanti L, Staats PS. Nampia-

- parampil DE, Hirsch JA. What is the role of epidural injections in the treatment of lumbar discogenic pain: A systematic review of comparative analysis with fusion and disc arthroplasty. *Korean J Pain* 2015; 28:75-87.
227. Manchikanti L, Falco FJE. Safeguards to prevent neurologic complications after epidural steroid injections: Analysis of evidence and lack of applicability of controversial policies. *Pain Physician* 2015; 18:E129-E138.
228. Manchikanti L, Hirsch JA. Neurological complications associated with epidural steroid injections. *Curr Pain Headache Rep* 2015; 19:482.
229. Manchikanti L, Pampati V, Benyamin RM, Boswell MV. Analysis of efficacy differences between caudal and lumbar interlaminar epidural injections in chronic lumbar axial discogenic pain: Local anesthetic alone vs. local combined with steroids. *Int J Med Sci* 2015; 12:214-222.
230. Manchikanti L, Singh V, Pampati V, Falco FJE, Hirsch JA. Comparison of the efficacy of caudal, interlaminar, and transforaminal epidural injections in managing lumbar disc herniation: Is one method superior to the other? *Korean J Pain* 2015; 28:11-21.
231. Manchikanti L, Falco FJE, Pampati V, Hirsch JA. Lumbar interlaminar epidural injections are superior to caudal epidural injections in managing lumbar central spinal stenosis. *Pain Physician* 2014; 17:E691-E702.
232. Manchikanti L, Falco FJE, Benyamin RM, Gharibo CG, Candido KD, Hirsch JA. Epidural steroid injections safety recommendations by the Multi-Society Pain Workgroup (MPW): More regulations without evidence or clarification. *Pain Physician* 2014; 17:E575-E588.
233. Manchikanti L, Candido KD, Singh V, Gharibo CG, Boswell MV, Benyamin RM, Falco FJE, Grider JS, Diwan S, Hirsch JA. Epidural steroid warning controversy still dogging FDA. *Pain Physician* 2014; 17:E451-E474.
234. Manchikanti L, Candido KD, Kaye AD, Boswell MV, Benyamin RM, Falco FJE, Gharibo CG, Hirsch JA. Randomized trial of epidural injections for spinal stenosis published in the New England Journal of Medicine: Further confusion without clarification. *Pain Physician* 2014; 17:E475-E488.
235. King W, Lau P, Lees R, Bogduk N. The validity of manual examination in assessing patients with neck pain. *Spine J* 2007; 7:22-26.
236. Carragee EJ, Haldeman S, Hurwitz E. The pyrite standard: The Midas touch in the diagnosis of axial pain syndromes. *Spine J* 2007; 7:27-31.
237. Bogduk N. In defense of King et al: The validity of manual examination in assessing patients with neck pain. *Spine J* 2007; 7:749-752.
238. Carragee EJ, Haldeman S, Hurwitz E. Response to Bogduk. Letter to the Editor. *Spine J* 2007; 7:752-753.
239. Bogduk N, McGuirk B. Causes and sources of chronic low back pain. In: Bogduk N, McGuirk B (eds). *Medical Management of Acute and Chronic Low Back Pain. An Evidence-Based Approach: Pain Research and Clinical Management*. Volume 13. Elsevier Science BV, Amsterdam, 2002, pp 115-126.
240. Barnsley L, Lord SM, Wallis BJ, Bogduk N. The prevalence of chronic cervical zygapophyseal joint pain after whiplash. *Spine (Phila Pa 1976)* 1995; 20:20-26.
241. Lord SM, Barnsley L, Wallis BJ, Bogduk N. Chronic cervical zygapophyseal joint pain with whiplash: A placebo-controlled prevalence study. *Spine (Phila Pa 1976)* 1996; 21:1737-1745.
242. Manchikanti L, Boswell MV, Singh V, Pampati V, Damron KS, Beyer CD. Prevalence of facet joint pain in chronic spinal pain of cervical, thoracic, and lumbar regions. *BMC Musculoskelet Disord* 2004; 5:15.
243. Manchikanti L, Pampati V, Damron KS, McManus CD, Jackson SD, Barnhill RC, Martin JC. A randomized, prospective, double-blind, placebo-controlled evaluation of the effect of sedation on diagnostic validity of cervical facet joint pain. *Pain Physician* 2004; 7:301-309.
244. Manchikanti L, Damron KS, Rivera J, McManus CD, Jackson SD, Barnhill RC, Martin JC. Evaluation of effect of sedation as a confounding factor in the diagnostic validity of lumbar facet joint pain: A prospective, randomized, double-blind, placebo-controlled evaluation. *Pain Physician* 2004; 7:411-417.
245. Barnsley L, Lord S, Wallis B, Bogduk N. False-positive rates of cervical zygapophyseal joint blocks. *Clin J Pain* 1993; 9:124-130.
246. Manchikanti L, Cash KA, Pampati V, Fellows B. Influence of psychological variables on the diagnosis of facet joint involvement in chronic spinal pain. *Pain Physician* 2008; 11:145-160.
247. Manchikanti L, Boswell MV, Manchukonda R, Cash KA, Giordano J. Influence of prior opioid exposure on diagnostic facet joint nerve blocks. *J Opioid Manage* 2008; 4:351-360.
248. Pampati S, Cash KA, Manchikanti L. Accuracy of diagnostic lumbar facet joint nerve blocks: A 2-year follow-up of 152 patients diagnosed with controlled diagnostic blocks. *Pain Physician* 2009; 12:855-866.
249. Manchikanti L, Pampati S, Cash KA. Making sense of the accuracy of diagnostic lumbar facet joint nerve blocks: An assessment of implications of 50% relief, 80% relief, single block or controlled diagnostic blocks. *Pain Physician* 2010; 13:133-143.
250. Saal JS. General principles of diagnostic testing as related to painful lumbar spine disorders. A critical appraisal of current diagnostic techniques. *Spine (Phila Pa 1976)* 2002; 27:2538-2545.